Index of Authors

ADAMS, R. L. P., HENDERSON, M. A. L., WOOD, W. & LINDSAY, J. G. Multiple forms of nuclear deoxyribonucleic acid polymerases and their relationship with the soluble enzyme 237–246
ADAMS, R. W. *see* JAMESON, G. W. 107–117
AINSWORTH, S. & MACFARLANE, N. A kinetic study of rabbit muscle pyruvate kinase 223–236
ALLEN, A. K., NEUBERGER, A. & SHARON, N. The purification, composition and specificity of wheat-germ agglutinin 155–162
ALLEN, G. A. *see* EGGERT, F. M. 471–484
AMBLER, R. P. & WYNN, M. The amino acid sequences of cytochromes *c*-551 from three species of *Pseudomonas* 485–498

BAILEY, A. J. *see* ROBINS, S. P. 771–780
BARCLAY, A. *see* CRICHTON, R. R. 855–857
BARDSLEY, W. G., CRABBE, M. J. C. & SHINDLER, J. S. Kinetics of the diamine oxidase reaction 459–469
BARNETT, J. E. G., HOLMAN, G. D. & MUNDAY, K. A. Structural requirements for binding to the sugar-transport system of the human erythrocyte 211–221
BARNETT, J. E. G., RASHEED, A. & CORINA, D. L. Partial reactions of D-glucose 6-phosphate–1L-*myo*inositol 1-phosphate cyclase 21–30
BARRETT, A. J. Human cathepsin B1. Purification and some properties of the enzyme 809–822
BARRETT, A. J. *see also* STARKEY, P. M. 823–831
BAR-TANA, J. & ROSE, G. Rat liver microsomal palmitoyl-coenzyme A synthetase. Structural properties 443–449
BAR-TANA, J., ROSE, G., BRANDES, R. & SHAPIRO, B. Palmitoyl-coenzyme A synthetase. Mechanism of reaction 199–209
BATTELLI, M. G., LORENZONI, E. & STIRPE, F. Milk xanthine oxidase type D (dehydrogenase) and type O (oxidase). Purification, interconversion and some properties 191–198
BERGERSEN, F. J. & TURNER, G. L. Kinetic studies of nitrogenase from soya-bean root-nodule bacteroids 61–75
BERNS, D. S. *see* KAO, O. H. W. 39–50
BESSELL, E. M. & THOMAS, P. The deoxyfluoro-D-glucopyranose 6-phosphates and their effect on yeast glucose phosphate isomerase 77–82
BESSELL, E. M. & THOMAS, P. The effect of substitution at C-2 of D-glucose 6-phosphate on the rate of dehydrogenation by glucose 6-phosphate dehydrogenase (from yeast and from rat liver) 83–89
BLACKBURN, G. M. *see* SPENCER, M. 421–423
BLACKBURN, P. *see* SPENCER, M. 421–423
BLASS, J. P. & LEWIS, C. A. Inhibition by acetaldehyde of the pyruvate dehydrogenase complex from ox brain and ox kidney 415–416
BLASS, J. P. & LEWIS, C. A. Kinetic properties of the partially purified pyruvate dehydrogenase complex of ox brain 31–37
BONAVENTURA, C. *see* WILSON, M. T. 863–865
BONAVENTURA, J. *see* WILSON, M. T. 863–865
BOSE, S. K. *see* SENGUPTA, S. 623–624
BOULTER, D. *see* BROWN, R. H. 247–251, 253–256
BOWEN, D. M. & DAVISON, A. N. Cathepsin A in human brain and spleen 417–419
BRADBURY, A. F. & SMYTH, D. G. The use of 3-bromopropionic acid for the determination of protein thiol groups 637–642
BRANDES, R. *see* BAR-TANA, J. 199–209
BROWN, P. R. *see* HUGHES, S. G. 583
BROWN, R. H. & BOULTER, D. The amino acid sequence of cytochrome *c* from *Allium porrum* L. (leek) 247–251
BROWN, R. H., RICHARDSON, M., SCOGIN, R. & BOULTER, D. The amino acid sequence of cytochrome *c* from *Spinacea oleracea* L. (spinach) 253–256
BROWNSON, C. & WATTS, D. C. The modification of cholinesterase activity by 5,5′-dithiobis-(2-nitrobenzoic acid) included in the coupled spectrophotometric assay. Evidence for a non-catalytic substrate-binding site 369–374
BRUNORI, M. *see* WILSON, M. T. 863–865
BRYCE, C. F. A. *see* CRICHTON, R. R. 51–59, 855–857
BURGESS, R. C. *see* EGGERT, F. M. 471–484
BUTTERWORTH, P. J. *see* RAMASAMY, I. 359–367
BYRDE, R. J. W. *see* REYES, F. 381–388

CARROLL, M. & ROBINSON, D. Immunological properties of *N*-acetyl-β-D-glucosaminidase of normal human liver and of GM_2-gangliosidosis liver 91–96
CHANCE, B. & OSHINO, N. Analysis of the catalase–hydrogen peroxide intermediate in coupled oxidations 564–567
CHANCE, B. *see also* OSHINO, N. 555–563
CHAPMAN, M. J., MILLS, G. L. & TAYLAUR, C. E. The effect of a lipid-rich diet on the properties and composition of lipoprotein particles from the Golgi apparatus of guinea-pig liver 177–185
CHASSEAUD, L. F. Distribution of enzymes that catalyse reactions of glutathione with αβ-unsaturated compounds 765–769
COLE, H. A. *see* PERRY, S. V. 425–428
CORINA, D. L. *see* BARNETT, J. E. G. 21–30
COULSON, A. *see* RENSING, U. F. E. 605–610
CRABBE, M. J. C. *see* BARDSLEY, W. G. 459–469
CRICHTON, R. R., EASON, R., BARCLAY, A. & BRYCE, C. F. A. The subunit structure of horse spleen apoferritin: the molecular weight of the oligomer and its stability to dissociation by dilution 855–857
CRICHTON, R. R., MILLAR, J. A., CUMMING, R. L. C. & BRYCE, C. F. A. The organ-specificity of ferritin in human and horse liver and spleen 51–59
CUMMING, R. L. C. *see* CRICHTON, R. R. 51–59

DAVIES, D. D. *see* WONG, K. F. 451–458
DAVISON, A. N. *see* BOWEN, D. M. 417–419
DE LA CHAPELLE, A. *see* SELANDER, R.-K. 757–764

DICKINSON, F. M. & MONGER, G. P. A study of the kinetics and mechanism of yeast alcohol dehydrogenase with a variety of substrates 261–270
DIXON, H. B. F. Shapes of curves of pH-dependence of reactions 149–154
DUKE, E. J., JOYCE, P. & RYAN, J. P. Characterization of alternative molecular forms of xanthine oxidase in the mouse 187–190
DUTTON, G. J. *see* ILLING, H. P. A. 139–147

EASON, R. *see* CRICHTON, R. R. 855–857
EGGERT, F. M., ALLEN, G. A. & BURGESS, R. C. Amelogenins. Purification and partial characterization of proteins from developing bovine dental enamel 471–484
ELDER, G. H. *see* STOLL, M. S. 429–432
ELMORE, D. T. *see* JAMESON, G. W. 107–117
ENGEL, P. C. & FERDINAND, W. The significance of abrupt transitions in Lineweaver–Burk plots with particular reference to glutamate dehydrogenase. Negative and positive co-operativity in catalytic rate constants 97–105
ERICKSON, J. S. *see* HO, M. W. 173–176
ESNOUF, M. P., LLOYD, P. H. & JESTY, J. A method for the simultaneous isolation of Factor X and prothrombin from bovine plasma 781–789
ESNOUF, M. P. *see also* JESTY, J. 791–797

FERDINAND, W. *see* ENGEL, P. C. 97–105; SPENCER, M. 421–423
FRÈRE, J.-M. *see* NIETO, M. 163–171; PERKINS, H. R. 707–718
FROSCIO, M. *see* KEMP, B. E. 271–274

GAMES, D. E. *see* STOLL, M. S. 429–432
GHUYSEN, J.-M. *see* NIETO, M. 163–171; PERKINS, H. R. 707–718
GREENAWAY, P. J. & MURRAY, K. Polynucleotides associated with histone preparations 585–591
GRIFFIN, D. C. *see* MAYES, R. W. 541–553

HARTLEY, B. S. *see* SHOTTON, D. M. 643–675
HATTON, M. W. C. Studies on the coagulant enzyme from *Agkistrodon rhodostoma* venom. Isolation and some properties of the enzyme 799–807
HAYTON, G. J., PEARSON, C. K., SCAIFE, J. R. & KEIR, H. M. Synthesis of deoxyribonucleic acid in BHK-21/C13 cells 499–508
HENDERSON, M. A. L. *see* ADAMS, R. L. P. 237–246
HO, M. W., O'BRIEN, J. S., RADIN, N. S. & ERICKSON, J. S. Glucocerebrosidase: reconstitution of activity from macromolecular components 173–176
HOLBROOK, J. J. & INGRAM, V. A. Ionic properties of an essential histidine residue in pig heart lactate dehydrogenase 729–738
HOLBROOK, J. J. & STINSON, R. A. The use of ternary complexes to study ionizations and isomerizations during catalysis by lactate dehydrogenase 739–748
HOLBROOK, J. J. *see also* STINSON, R. A. 719–728
HOLMAN, G. D. *see* BARNETT, J. E. G. 211–221

HUGHES, S. G. & BROWN, P. R. Improved preparation of bacteriophage T4-infected *Escherichia coli* for the isolation of polynucleotide kinase 583
HUNT, J. A. Interaction between polyuridylic acid and rabbit globin messenger ribonucleic acid 327–333
HUNT, J. A. Terminal sequence studies of high-molecular-weight ribonucleic acid. The 3′ termini of rabbit globin messenger ribonucleic acid 315–325

ILLING, H. P. A. & DUTTON, G. J. Some properties of the uridine diphosphate glucuronyltransferase activity synthesizing thio-β-D-glucuronides 139–147
INGLE, J. *see* JACKSON, M. 523–533
INGRAM, V. A. *see* HOLBROOK, J. J. 729–738
INMAN, F. P. *see* RICARDO, M. J., JR. 677–682
IRVING, M. G. & WILLIAMS, J. F. Kinetic studies on the regulation of rabbit liver pyruvate kinase 287–301
IRVING, M. G. & WILLIAMS, J. F. Studies on the interaction between rabbit liver pyruvate kinase and its allosteric effector fructose 1,6-diphosphate 303–313
ISEMURA, M., ZAHN, R. K. & SCHMID, K. A new neuraminic acid derivative and three types of glycopeptides isolated from the Cuvierian tubules of the sea cucumber *Holothuria forskali* 509–521

JACKSON, A. H. *see* STOLL, M. S. 429–432
JACKSON, M. & INGLE, J. The nature of polydisperse ribonucleic acid in plants 523–533
JAMESON, G. W., ROBERTS, D. V., ADAMS, R. W., KYLE, W. S. A. & ELMORE, D. T. Determination of the operational molarity of solutions of bovine α-chymotrypsin, trypsin, thrombin and Factor X*a* by spectrofluorimetric titration 107–117
JESTY, J. & ESNOUF, M. P. The preparation of activated Factor X and its action on prothrombin 791–797
JESTY, J. *see also* ESNOUF, M. P. 781–789
JOYCE, P. *see* DUKE, E. J. 187–190

KAO, O. H. W., BERNS, D. S. & TOWN, W. R. The characterization of C-phycocyanin from an extremely halo-tolerant blue-green alga, *Coccochloris elabens* 39–50
KEIR, H. M. *see* HAYTON, G. J. 499–508
KEMP, B. E., FROSCIO, M. & MURRAY, A. W. Protein kinase activity in commercially available crystalline yeast alcohol dehydrogenase 271–274
KIDO, R. *see* NOGUCHI, T. 375–380
KNOWLER, J. T. & SMELLIE, R. M. S. The oestrogen-stimulated synthesis of heterogeneous nuclear ribonucleic acid in the uterus of immature rats 689–697
KRISHNAN, P. S. *see* SREE KUMAR, K. 683–687
KYLE, W. S. A. *see* JAMESON, G. W. 107–117

LAPETINA, E. G. & MICHELL, R. H. A membrane-bound activity catalysing phosphatidylinositol breakdown to 1,2-diacylglycerol, D-*myo*inositol 1:2-cyclic phosphate and D-*myo*inositol 1-phosphate. Properties and subcellular distribution in rat cerebral cortex 433–442
LEWIS, C. A. *see* BLASS, J. P. 31–37, 415–416
LEYH-BOUILLE, M. *see* NIETO, M. 163–171; PERKINS, H. R. 707–718

LINDSAY, J. G. *see* ADAMS, R. L. P. 237–246
LLOYD, P. H. *see* ESNOUF, M. P. 781–789
LORENZONI, E. *see* BATTELLI, M. G. 191–198

MACFARLANE, N. *see* AINSWORTH, S. 223–236
MAK, D. *see* ROLLESTON, F. S. 851–853
MASON, R. M. & MAYES, R. W. Extraction of cartilage protein–polysaccharides with inorganic salt solutions 535–540
MASON, R. M. *see also* MAYES, R. W. 541–553
MAYES, R. W., MASON, R. M. & GRIFFIN, D. C. The composition of cartilage proteoglycans. An investigation using high- and low-ionic-strength extraction procedures 541–553
MAYES, R. W. *see also* MASON, R. M. 535–540
MICHELL, R. H. *see* LAPETINA, E. G. 433–442
MILLAR, J. A. *see* CRICHTON, R. R. 51–59
MILLINGTON, D. S. *see* STOLL, M. S. 429–432
MILLS, G. L. *see* CHAPMAN, M. J. 177–185
MONGER, G. P. *see* DICKINSON, F. M. 261–270
MUNDAY, K. A. *see* BARNETT, J. E. G. 211–221
MURRAY, A. W. *see* KEMP, B. E. 271–274
MURRAY, K. Nucleotide sequence analysis with polynucleotide kinase and nucleotide 'mapping' methods. 5′-Terminal sequences of deoxyribonucleic acid from bacteriophages λ and 424 569–583
MURRAY, K. *see also* GREENAWAY, P. J. 585–591

NEUBERGER, A. *see* ALLEN, A. K. 155–162
NIETO, M., PERKINS, H. R., LEYH-BOUILLE, M., FRÈRE, J.-M. & GHUYSEN, J.-M. Peptide inhibitors of *Streptomyces* DD-carboxypeptidases 163–171
NIETO, M. *see also* PERKINS, H. R. 707–718
NISHINO, M. *see* NOGUCHI, T. 375–380
NOGUCHI, T., NISHINO, M. & KIDO, R. Tryptophan 5-hydroxylase in rat intestine 375–380

O'BRIEN, J. S. *see* HO, M. W. 173–176
OGSTON, A. G. & PRESTON, B. N. A sensitive and accurate gel osmometer 843–850
O'HANLON, P. *see* STOLL, M. S. 429–432
OSBORNE, D. J. *see* ROBERTS, B. E. 275–286
OSHINO, N., OSHINO, R. & CHANCE, B. The characteristics of the 'peroxidatic' reaction of catalase in ethanol oxidation 555–563
OSHINO, N. *see also* CHANCE, B. 564–567
OSHINO, R. *see* OSHINO, N. 555–563

PARKER, J. E. *see* TRIM, A. R. 257–259
PAVLIK, J. G. & ROGERS, H. J. Selective extraction of polymers from cell walls of Gram-positive bacteria 619–621
PAYNE, P. I. *see* ROBERTS, B. E. 275–286
PEARSON, C. K. *see* HAYTON, G. J. 499–508
PERHAM, R. N. The reactivity of functional groups as a probe for investigating the topography of tobacco mosaic virus. The use of mutants with additional lysine residues in the coat protein 119–126
PERHAM, R. N. *see also* RILEY, M. 625–635; STRIBLING, D. 833–841
PERKINS, H. R., NIETO, M., FRÈRE, J.-M., LEYH-BOUILLE, M. & GHUYSEN, J.-M. *Streptomyces* DD-carboxypeptidases as transpeptidases. The specificity for amino compounds acting as carboxyl acceptors 707–718
PERKINS, H. R. *see also* NIETO, M. 163–171
PERRY, S. V. & COLE, H. A. Phosphorylation of the '37000 component' of the troponin complex (troponin-T) 425–428
PERRY, S. V. *see also* STONE, D. 127–137
PRESTON, B. N. *see* OGSTON, A. G. 843–850

RADIN, N. S. *see* HO, M. W. 173–176
RAMASAMY, I. & BUTTERWORTH, P. J. The inhibition of pig kidney alkaline phosphatase by oxidized or reduced nicotinamide–adenine dinucleotide and related compounds 359–367
RASHEED, A. *see* BARNETT, J. E. G. 21–30
RENSING, U. F. E. A sequence of seventy-three nucleotides from the coliphage R17 genome 593–604
RENSING, U. F. E., COULSON, A. & THOMPSON, E. O. P. Nucleotide sequences of similar size from the coliphage R17 genome 605–610
REYES, F. & BYRDE, R. J. W. Partial purification and properties of a β-*N*-acetylglucosaminidase from the fungus *Sclerotinia fructigena* 381–388
RICARDO, M. J., JR. & INMAN, F. P. Investigations of the structural function of the J chain in human immunoglobulin M 677–682
RICHARDSON, M. *see* BROWN, R. H. 253–256
RILEY, M. & PERHAM, R. N. The reaction of protein amino groups with methyl 5-iodopyridine-2-carboximidate. A possible general method of preparing isomorphous heavy-atom derivatives of proteins 625–635
ROBERTS, B. E., PAYNE, P. I. & OSBORNE, D. J. Protein synthesis and the viability of rye grains. Loss of activity of protein-synthesizing systems *in vitro* associated with a loss of viability 275–286
ROBERTS, D. V. *see* JAMESON, G. W. 107–117
ROBINS, S. P., SHIMOKOMAKI, M. & BAILEY, A. J. The chemistry of the collagen cross-links. Age-related changes in the reducible components of intact bovine collagen fibres 771–780
ROBINSON, D. *see* CARROLL, M. 91–96
ROGERS, H. J. *see* PAVLIK, J. G. 619–621
ROLLESTON, F. S. & MAK, D. The binding of polyribosomes to smooth and rough endoplasmic-reticulum membranes 851–853
ROSE, G. *see* BAR-TANA, J. 199–209, 443–449
RUSSELL, A. E. Effect of alcohols and neutral salt on the thermal stability of soluble and precipitated acid-soluble collagen 335–342
RYAN, J. P. *see* DUKE, E. J. 187–190

SCAIFE, J. R. *see* HAYTON, G. J. 499–508
SCHMID, K. *see* ISEMURA, M. 509–521
SCOGIN, R. *see* BROWN, R. H. 253–256
SELANDER, R.-K. Interaction of quinacrine mustard with mononucleotides and polynucleotides 749–755
SELANDER, R.-K. & DE LA CHAPELLE, A. Interaction of quinacrine mustard with calf thymus histone fractions 757–764

SENGUPTA, S. & BOSE, S. K. Stereoconfiguration of amino acids in peptides from a mycobacillin-synthesizing cell-free system 623–624
SHAPIRO, B. *see* BAR-TANA, J. 199–209
SHARON, N. *see* ALLEN, A. K. 155–162
SHIMOKOMAKI, M. *see* ROBINS, S. P. 771–780
SHINDLER, J. S. *see* BARDSLEY, W. G. 459–469
SHOTTON, D. M. & HARTLEY, B. S. Evidence for the amino acid sequence of porcine pancreatic elastase 643–675
SILVER, M. J. *see* SMITH, J. B. 615–618
SIMS, P. Epoxy derivatives of aromatic polycyclic hydrocarbons. The preparation and metabolism of epoxides related to 7,12-dimethylbenz[*a*]anthracene 405–413
SITARAMAYYA, A. *see* SREE KUMAR, K. 683–687
SMELLIE, R. M. S. *see* KNOWLER, J. T. 689–697
SMITH, J. A. & STOCKEN, L. A. The characterization of a non-histone protein isolated from histone F1 preparations 859–861
SMITH, J. B., SILVER, M. J. & WEBSTER, G. R. Phospholipase A_1 of human blood platelets 615–618
SMYTH, D. G. *see* BRADBURY, A. F. 637–642
SPENCER, M., BLACKBURN, P., FERDINAND, W. & BLACKBURN, G. M. Competitive inhibition and substrate activity of uridine diphosphate 6-deoxygalactose for *Escherichia coli* uridine diphosphate galactose 4-epimerase 421–423
SREE KUMAR, K., SITARAMAYYA, A. & KRISHNAN, P. S. Modulation of guanine deaminase 683–687
STARKEY, P. M. & BARRETT, A. J. Human cathepsin B1. Inhibition by α_2-macroglobulin and other serum proteins 823–831
STINSON, R. A. & HOLBROOK, J. J. Equilibrium binding of nicotinamide nucleotides to lactate dehydrogenases 719–728
STINSON, R. A. *see also* HOLBROOK, J. J. 739–748
STIRPE, F. *see* BATTELLI, M. G. 191–198
STOCKEN, L. A. *see* SMITH, J. A. 859–861
STOLL, M. S., ELDER, G. H., GAMES, D. E., O'HANLON, P., MILLINGTON, D. S. & JACKSON, A. H. Isocoproporphyrin: nuclear-magnetic-resonance- and mass-spectral methods for the determination of porphyrin structure 429–432
STONE, D. & PERRY, S. V. Studies on the heterogeneity of subfragment-1 preparations. Isolation of a new proteolytic fragment of the heavy chain of myosin 127–137
STONE, K. J. & TOWNSLEY, B. H. The effect of L-ascorbate on catecholamine biosynthesis 611–613
STRIBLING, D. & PERHAM, R. N. Purification and characterization of two fructose diphosphate aldolases from *Escherichia coli* (Crookes' strain) 833–841

TAIT, G. H. Aminolaevulinate synthetase of *Micrococcus denitrificans*. Purification and properties of the enzyme, and the effect of growth conditions on the enzyme activity in cells 389–403
TAYLAUR, C. E. *see* CHAPMAN, M. J. 177–185
THOMAS, P. *see* BESSELL, E. M. 77–82, 83–89
THOMPSON, E. O. P. *see* RENSING, U. F. E. 605–610
TOWN, W. R. *see* KAO, O. H. W. 39–50
TOWNSLEY, B. H. *see* STONE, K. J. 611–613
TRIM, A. R. & PARKER, J. E. The preparation and analysis of a new trinucleotide from yeast ribonucleic acid, containing two adjacent 2′-*O*-methylpentose residues 257–259
TURNER, G. L. *see* BERGERSEN, F. J. 61–75

WATTS, D. C. *see* BROWNSON, C. 369–374
WEBSTER, G. R. *see* SMITH, J. B. 615–618
WHITEMAN, P. The quantitative determination of glycosaminoglycans in urine with Alcian Blue 8GX 351–357
WHITEMAN, P. The quantitative measurement of Alcian Blue–glycosaminoglycan complexes 343–350
WILLIAMS, J. F. *see* IRVING, M. G. 287–301, 303–313
WILSON, M. T., BRUNORI, M., BONAVENTURA, J. & BONAVENTURA, C. Effect of steady illumination on the binding of carbon monoxide by carboxymethylated cytochrome *c* 863–865
WONG, K. F. & DAVIES, D. D. Regulation of phosphoenolpyruvate carboxylase of *Zea mays* by metabolites 451–458
WOOD, W. *see* ADAMS, R. L. P. 237–246
WYNN, M. *see* AMBLER, R. P. 485–498

YON, R. J. Wheat-germ aspartate transcarbamoylase. The effects of ligands on the inactivation of the enzyme by trypsin and denaturing agents 699–706

ZAHN, R. K. *see* ISEMURA, M. 509–521

Index of Subjects

Acetaldehyde, inhibition by, of the pyruvate dehydrogenase complex from ox cerebral cortex and ox kidney (Blass, J. P. & Lewis, C. A.) 415–416

Acetaldehyde, kinetics and mechanism of the reaction catalysed by baker's-yeast alcohol dehydrogenase with, and a variety of other substrates (Dickinson, F. M. & Monger, G. P.) 261–270

Acetylcholinesterase, erythrocyte, human, modification of the activity of, by 5,5′-dithiobis-(2-nitrobenzoate) included in the coupled spectrophotometric assay (Brownson, C. & Watts, D. C.) 369–374

Acetyl-coenzyme A, lack of effect of, on partially purified ox cerebral-cortex pyruvate dehydrogenase complex (Blass, J. P. & Lewis, C. A.) 31–37

Acetylene, kinetics of the reduction of nitrogen and, by the nitrogenase complex from soya-bean root-nodule bacteroids (Bergersen, F. J. & Turner, G. L.) 61–75

N-Acetylglucosamine, inhibition by, of wheat-germ agglutinin (Allen, A. K., Neuberger, A. & Sharon, N.) 155–162

N-Acetyl-β-D-glucosaminidase, liver, immunological properties of, from normal humans and from patients with GM_2-gangliosidosis (Carroll, M. & Robinson, D.) 91–96

β-*N*-Acetylglucosaminidase, partial purification and properties of, from *Sclerotinia fructigena* (Reyes, F. & Byrde, R. J. W.) 381–388

Acid β-glucosidase, *see* β-Glucosidase, acid

Adenine nucleotides, effects of, on the activity of maize-seedling phosphoenolpyruvate carboxylase (Wong, K. F. & Davies, D. D.) 451–458

Adenosine diphosphate, kinetics of the reaction between phosphoenolpyruvate and, catalysed by rabbit skeletal-muscle pyruvate kinase (Ainsworth, S. & Macfarlane, N.) 223–236

Adenosine, inhibition by, of pig kidney alkaline phosphatase (Ramasamy, I. & Butterworth, P. J.) 359–367

Adenosine triphosphatase, activity of, of subfragment 1 preparations obtained by proteolysis with papain from the heavy chain of rabbit skeletal-muscle myosin (Stone, D. & Perry, S. V.) 127–137

Adenosine triphosphate, inhibition by, of the activity of rabbit liver pyruvate kinase (Irving, M. G. & Williams, J. F.) 287–301

Agglutinin, wheat-germ, purification, composition and specificity of (Allen, A. K., Neuberger, A. & Sharon, N.) 155–162

Agkistrodon rhodostoma, *see* Malayan pit viper

Alanine, inhibition by, of the activity of rabbit liver pyruvate kinase (Irving, M. G. & Williams, J. F.) 287–301

Albumin, plasma, ox, interaction of quinacrine mustard with, and other proteins (Selander, R.-K. & de la Chapelle, A.) 757–764

Alcian Blue 8GX, quantitative determination of glycosaminoglycans in urine with (Whiteman, P.) 351–357

Alcian Blue 8GX, quantitative measurement of complexes of glycosaminoglycans with (Whiteman, P.) 343–350

Alcohol dehydrogenase, baker's-yeast, kinetics and mechanism of the reaction catalysed by, with a variety of substrates (Dickinson, F. M. & Monger, G. P.) 261–270

Alcohol dehydrogenase, yeast, protein kinase activity in commercially available crystalline preparations of (Kemp, B. E., Froscio, M. & Murray, A. W.) 271–274

Alcohols, effect of neutral salt and, on the thermal stability of soluble and precipitated acid-soluble calf skin collagen (Russell, A. E.) 335–342

Aldolase, *see* Fructose diphosphate aldolase

Alkaline phosphatase, *see* Phosphatase, alkaline

Allantoin, effects of, on the activities of rat and mouse liver and brain guanine deaminase isoenzymes (Sree Kumar, K., Sitaramayya, A. & Krishnan, P. S.) 683–687

Allium porrum, *see* Leek

Amelogenins, purification and partial characterization of, from developing ox dental enamel (Eggert, F. M., Allen, G. A. & Burgess, R. C.) 471–484

Amino acid compositions of amelogenins purified from developing ox dental enamel (Eggert, F. M., Allen, G. A. & Burgess, R. C.) 471–484

Amino acid residues, stereoconfiguration of, in peptides isolated from a mycobacillin-synthesizing cell-free system prepared from *Bacillus subtilis* B_3 (Sengupta, S. & Bose, S. K.) 623–624

Amino acid sequence, complete, evidence for, of pig pancreas elastase (Shotton, D. M. & Hartley, B. S.) 643–675

Amino acid sequence of cytochrome *c* from leek (Brown, R. H. & Boulter, D.) 247–251

Amino acid sequence of cytochrome *c* from spinach (Brown, R. H., Richardson, M., Scogin, R. & Boulter, D.) 253–256

Amino acid sequences of cytochromes *c*-551 from *Pseudomonas fluorescens* A.T.C.C. 17400, *Pseudomonas stutzeri* A.T.C.C. 17588 and *Pseudomonas mendocina* (Ambler, R. P. & Wynn, M.) 485–498

Amino compounds, specificity of *Streptomyces* DD-carboxypeptidases acting as peptidyltransferases for, as carboxyl acceptors (Perkins, H. R., Nieto, M., Frère, J.-M., Leyh-Bouille, M. & Ghuysen, J.-M.) 707–718

Amino groups, protein, reaction of methyl iodopicolinimidate with, and its possible use as a general method for preparing isomorphous heavy-atom derivatives of proteins (Riley, M. & Perham, R. N.) 625–635

5-Aminolaevulinate synthetase, purification and properties of, from *Micrococcus denitrificans* (Tait, G. H.) 389–403

3-Amino-L-tyrosine, reaction of methyl iodopicolinimidate with (Riley, M. & Perham, R. N.) 625–635

Apoferritin, organ-specificity of, in human and horse liver and spleen (Crichton, R. R., Millar, J. A., Cumming, R. L. C. & Bryce, C. F. A.) 51–59

Apoferritin, spleen, horse, molecular weight of the oligomer of, and its stability to dissociation by dilution (Crichton, R. R., Eason, R., Barclay, A. & Bruce, C. F. A.) 855–857

Arginine esterase, correlation of the coagulant activity of Arvin from the venom of the Malayan pit viper with activity of (Hatton, M. W. C.) 799–807
Arvin, isolation and properties of, from the venom of the Malayan pit viper (Hatton, M. W. C.) 799–807
L-Ascorbate, stimulation by, of the activity of rat liver phenylalanine hydroxylase (Stone, K. J. & Townsley, B. H.) 611–613
Aspartate carbamoyltransferase, wheat-germ, effects of ligands on the inactivation of, by trypsin and denaturing agents (Yon, R. J.) 699–706
Aspartate, inhibition by, of maize-seedling phosphoenolpyruvate carboxylase (Wong, K. F. & Davies, D. D.) 451–458

Bacillus licheniformis N.C.T.C. 6346, selective extraction of teichuronic acids from the cell walls of, and other Gram-positive bacteria (Pavlik, J. G. & Rogers, H. J.) 619–621
Bacillus subtilis B_3, stereoconfiguration of the amino acid residues in peptides isolated from a mycobacillin-synthesizing cell-free system prepared from (Sengupta, S. & Bose, S. K.) 623–624
Bacillus subtilis 168, selective extraction of teichuronic acids from the cell walls of, and other Gram-positive bacteria (Pavilk, J. G. & Rogers, H. J.) 619–621
Bacteriophage T4, improved preparation of *Escherichia coli* infected with, for the isolation of polynucleotide kinase (Hughes, S. G. & Brown, P. R.) 583
Bacteriophage λ, use of polynucleotide kinase and nucleotide 'mapping' methods for the determination of the 5′-hydroxy-terminal nucleotide sequences of deoxyribonucleic acid from (Murray, K.) 569–582
Bacteriophage 424, use of polynucleotide kinase and nucleotide 'mapping' methods for the determination of the 5′-hydroxy-terminal nucleotide sequences of deoxyribonucleic acid from (Murray, K.) 569–582
Bacteroids, root-nodule, soya-bean, kinetic properties of the nitrogenase complex from (Bergersen, F. J. & Turner, G. L.) 61–75
trans-Benzylideneacetone, distribution of enzymes that catalyse reactions of reduced glutathione with, and other $\alpha\beta$-unsaturated compounds in the livers of various vertebrate species (Chasseaud, L. F.) 765–769
Blood, human, modification of the activities of acetylcholinesterase and cholinesterase from, by 5,5′-dithiobis-(2-nitrobenzoate) included in the coupled spectrophotometric assay (Brownson, C. & Watts, D. C.) 369–374
Blood platelets, human, activity of phospholipase A_1 in (Smith, J. B., Silver, M. J. & Webster, G. R.) 615–618
Brain cortex, human, activity of cathepsin A in (Bowen, D. M. & Davison, A. N.) 417–419
Brain cortex, ox, inhibition by acetaldehyde of the pyruvate dehydrogenase complex from (Blass, J. P. & Lewis, C. A.) 415–416
Brain cortex, ox, kinetic properties of the partially purified pyruvate dehydrogenase complex from (Blass, J. P. & Lewis, C. A.) 31–37
Brain cortex, rat, properties and subcellular distribution of a membrane-bound enzyme activity catalysing the breakdown of phosphatidylinositol to 1,2-diacylglycerol, D-*myo*inositol 1:2-cyclic phosphate and D-*myo*inositol 1-phosphate in (Lapetina, E. G. & Michell, R. H.) 433–442
Brain, rat and mouse, regulation of the activities of guanine deaminase isoenzymes in rat and mouse liver and (Sree Kumar, K., Sitaramayya, A. & Krishnan, P. S.) 683–687
3-Bromopropionic acid, use of, for the determination of protein thiol groups (Bradbury, A. F. & Smyth, D. G.) 637–642
Butan-1-ol, kinetics and mechanism of the reaction catalysed by baker's-yeast alcohol dehydrogenase with, and a variety of other substrates (Dickinson, F. M. & Monger, G. P.) 261–270
Butyraldehyde, kinetics and mechanism of the reaction catalysed by baker's-yeast alcohol dehydrogenase with, and a variety of other substrates (Dickinson, F. M. & Monger, G. P.) 261–270

Calcium ions, effect of, on anaerobic growth and 5-aminolaevulinate synthetase activity of *Micrococcus denitrificans* (Tait, G. H.) 389–403
Carbamoyl phosphate, antagonization by, of the protecting effect of uridine monophosphate against inactivation of wheat-germ aspartate carbamoyltransferase and denaturing agents (Yon, R. J.) 699–706
Carbohydrate compositions of three types of glycopeptides isolated from the Cuvierian tubules of the sea-cucumber *Holothuria forskali* (Isemura, M., Zahn, R. K. & Schmid, K.) 509–521
Carbon monoxide, effect of steady illumination on the binding of, by carboxymethylated horse heart cytochrome *c* (Wilson, M. T., Brunori, M., Bonaventura, J. & Bonaventura, C.) 863–865
S-Carboxyethylcysteine, formation of, in the reaction of 3-bromopropionic acid with cysteine residues in proteins (Bradbury, A. F. & Smyth, D. G.) 637–642
DD-Carboxypeptidases, peptide inhibitors of, from *Streptomyces* strains (Nieto, M., Perkins, H. R., Leyh-Bouille, M., Frère, J.-M. & Ghuysen, J.-M.) 163–171
DD-Carboxypeptidases, *Streptomyces*, specificity of, acting as peptidyltransferases for amino compounds as carboxyl acceptors (Perkins, H. R., Nieto, M., Frère, J.-M., Leyh-Bouille, M. & Ghuysen, J.-M.) 707–718
Cartilage, articular, ox, age-related changes in the reducible cross-links of intact collagen fibres from (Robins, S. P., Shimokomaki, M. & Bailey, A. J.) 771–780
Cartilage, nasal, ox, composition of proteoglycans extracted from, with solutions of high and of low ionic strengths (Mayes, R. W., Mason, R. M. & Griffin, D. C.) 541–553
Cartilage, nasal, ox, extraction of proteoglycans from, with inorganic salt solutions of various ionic strengths (Mason, R. M. & Mayes, R. W.) 535–540
Catalase–hydrogen peroxide intermediate, kinetic analysis of, in coupled oxidations (Chance, B. & Oshino, N.) 564–567
Catalase, liver, rat, characteristics of the 'peroxidatic' reaction of, in the oxidation of ethanol (Oshino, N., Oshino, R. & Chance, B.) 555–563

Catecholamines, stimulation by ascorbate of the activity of rat liver phenylalanine hydroxylase and its possible role in the regulation of the biosynthesis of (Stone, K. J. & Townsley, B. H.) 611–613
Cathepsin A, activity of, in human cerebral cortex and spleen (Bowen, D. M. & Davison, A. N.) 417–419
Cathepsin B1, liver, human, inhibition by α_2-macroglobulin and other human serum proteins of the activity of (Starkey, P. M. & Barrett, A. J.) 823–831
Cathepsin B1, purification and properties of, from human liver (Barrett, A. J.) 809–821
Cell walls, selective extraction of teichuronic acids from, of Gram-positive bacteria (Pavlik, J. G. & Rogers, H. J.) 619–621
Cells, kidney, baby-hamster (BHK-21/C13), biosynthesis of deoxyribonucleic acid in (Hayton, G. J., Pearson, C. K., Scaife, J. R. & Keir, H. M.) 499–508
Cells, L929, mouse, cultured, multiple forms of nuclear deoxyribonucleic acid polymerases and their relationship with the cytosol enzyme from (Adams, R. L. P., Henderson, M. A. L., Wood, W. & Lindsay, J. G.) 237–246
Cerebral cortex, human, activity of cathepsin A in (Bowen, D. M. & Davison, A. N.) 417–419
Cerebral cortex, ox, inhibition by acetaldehyde of the pyruvate dehydrogenase complex from (Blass, J. P. & Lewis, C. A.) 415–416
Cerebral cortex, ox, kinetic properties of the partially purified pyruvate dehydrogenase complex from (Blass, J. P. & Lewis, C. A.) 31–37
Cerebral cortex, rat, properties and subcellular distribution of a membrane-bound enzyme activity catalysing the breakdown of phosphatidylinositol to 1,2-diacylglycerol, D-*myo*inositol 1:2-cyclic phosphate and D-*myo*inositol 1-phosphate in (Lapetina, E. G. & Michell, R. H.) 433–442
Chick embryo, properties of the uridine diphosphate glucuronyltransferase activity synthesizing thio-β-D-glucuronides in liver preparations from (Illing, H. P. A. & Dutton, G. J.) 139–147
Chloride, potassium, neutral, effect of alcohols and, on the thermal stability of soluble and precipitated acid-soluble calf skin collagen (Russell, A. E.) 335–342
o-Chlorobenzylidenemalononitrile, distribution of enzymes that catalyse reactions of reduced glutathione with, and other '$\alpha\beta$-unsaturated compounds in the livers of various vertebrate species (Chasseaud, L. F.) 765–769
Cholinesterase, plasma, human, modification of the activity of, by 5,5′-dithiobis-(2-nitrobenzoate) included in the coupled spectrophotometric assay (Brownson, C. & Watts, D. C.) 369–374
Chondroitin sulphate proteolgycan, extraction of, from ox nasal cartilage with solutions of low ionic strength (Mayes, R. W., Mason, R. M. & Griffin, D. C.) 541–553
Chromatin, properties of polynucleotides associated with preparations of histones obtained from, from various sources (Greenaway, P. J. & Murray, K.) 585–591
α-Chymotrypsin, ox, determination of the operational molarity of solutions of, by spectrofluorimetric titration (Jameson, G. W., Roberts, D. V., Adams, R. W., Kyle, W. S. A. & Elmore, D. T.) 107–117
Cinnamonitrile, distribution of enzymes that catalyse reactions of reduced glutathione with, and other $\alpha\beta$-unsaturated compounds in the livers of various vertebrate species (Chasseaud, L. F.) 765–769
Clotting Factor X, method for the simultaneous isolation of prothrombin and, from ox plasma (Esnouf, M. P., Lloyd, P. H. & Jesty, J.) 781–789
Clotting Factor X, plasma, ox, preparation of Factor X*a* from, and its action on prothrombin (Jesty, J. & Esnouf, M. P.) 791–797
Clotting Factor X*a*, preparation of, from ox plasma Factor X and its action on prothrombin (Jesty, J. & Esnouf, M. P.) 791–797
Coagulant enzyme (Arvin), isolation and properties of, from the venom of the Malayan pit viper (Hatton, M. W. C.) 799–807
Coat protein, reactivity towards methyl picolinimidate of additional lysine residues in, of mutants of tobacco-mosaic virus and the reagent's usefulness as a probe for investigating the topography of the virus (Perham, R. N.) 119–126
Coccochloris elabens, characterization of C-phycocyanin from (Kao, O. H. W., Berns, D. S. & Town, W. R.) 39–50
Coliphage R17, nucleotide sequence containing seventy-three residues from the genome of (Rensing, U. F. E.) 593–604
Coliphage R17, nucleotide sequences of similar size from the genome of (Rensing, U. F. E., Coulson, A. & Thompson, E. O. P.) 605–610
Collagen, age-related changes in the reducible cross-links of intact fibres of, from ox skin, tendon and articular cartilage (Robins, S. P., Shimokomaki, M. & Bailey, A. J.) 771–780
Collagen, skin, calf, acid-soluble, soluble and precipitated, effect of alcohols and neutral salt on the thermal stability of (Russell, A. E.) 335–342
Cupric ions, effect of, on anaerobic growth and 5-aminolaevulinate synthetase activity of *Micrococcus denitrificans* (Tait, G. H.) 389–403
Cuvierian tubules, sea-cucumber (*Holothuria forskali*), isolation of a new neuraminic acid derivative and three types of glycopeptides from (Isemura, M., Zahn, R. K. & Schmid, K.) 509–521
D-Cycloserine residues, inhibition by peptides containing, of DD-carboxypeptidases from *Streptomyces* strains (Nieto, M., Perkins, H. R., Leyh-Bouille, M., Frère, J.-M. & Ghuysen, J.-M.) 163–171
D-Cycloserine, specificity of *Streptomyces* DD-carboxypeptidases acting as peptidyltransferases for, and other amino compounds as carboxyl acceptors (Perkins, H. R., Nieto, M., Frère, J.-M., Leyh-Bouille, M. & Ghuysen, J.-M.) 707–718
Cysteine residues, reaction of 3-bromopropionic acid with, in proteins (Bradbury, A. F. & Smyth, D. G.) 637–642
Cytochrome *c*, heart, horse, accurate determination of the molecular weight of, with a sensitive polyacrylamide-gel osmometer (Ogston, A. G. & Preston, B. N.) 843–850
Cytochrome *c*, heart, horse, carboxymethylated, effect of steady illumination on the binding of carbon monoxide by (Wilson, M. T., Brunori, M., Bonaventura, J. & Bonaventura, C.) 863–865

Cytochrome *c*, leek, amino acid sequence of (Brown, R. H. & Boulter, D.) 247–251
Cytochrome *c*, spinach, amino acid sequence of (Brown, R. H., Richardson, M., Scogin, R. & Boulter, D.) 253–256
Cytochromes *c*-551, amino acid sequences of, from *Pseudomonas fluorescens* A.T.C.C. 17400, *Pseudomonas stutzeri* A.T.C.C. 17588 and *Pseudomonas mendocina* (Ambler, R. P. & Wynn, M.) 485–498
Cytosol, multiple forms of nuclear deoxyribonucleic acid polymerases and their relationship with the enzyme from, of cultured mouse L929 cells (Adams, R. L. P., Henderson, M. A. L., Wood, W. & Lindsay, J. G.) 237–246

Dental enamel, *see* Enamel, dental
Deoxyfluoro-D-glucose 6-phosphates, activity of yeast and rat liver glucose 6-phosphate dehydrogenase towards (Bessell, E. M. & Thomas, P.) 83–89
Deoxyfluoro-D-glucose 6-phosphates, activity of yeast glucose phosphate isomerase towards (Bessell, E. M. & Thomas, P.) 77–82
Deoxyribonucleic acid, biosynthesis of, in baby-hamster kidney (BHK-21/C13) cells (Hayton, G. J., Pearson, C. K., Scaife, J. R. & Keir, H. M.) 499–508
Deoxyribonucleic acid polymerases, nuclear, multiple forms of, and their relationship with the cytosol enzyme from cultured mouse L929 cells (Adams, R. L. P., Henderson, M. A. L., Wood, W. & Lindsay, J. G.) 237–246
Deoxyribonucleic acids, interaction of quinacrine mustard with (Selander, R.-K.) 749–755
Deoxyribonucleic acids, use of polynucleotide kinase and nucleotide 'mapping' methods for the determination of the 5′-hydroxy-terminal nucleotide sequences of, from bacteriophages λ and 424 (Murray, K.) 569–582
Deoxyribonucleoproteins, properties of polynucleotides associated with preparations of histones obtained from, from various sources (Greenaway, P. J. & Murray, K.) 585–591
Deuterium isotope (^{2}H), effect of the presence of, on the kinetics of the oxidation of *p*-dimethylaminomethyl[$\alpha\alpha$-$^2H_2$]benzylamine by pig kidney diamine oxidase (Bardsley, W. G., Crabbe, M. J. C. & Shindler, J. S.) 459–469
Development, changes in the reducible cross-links of intact collagen fibres from ox skin, tendon and articular cartilage during (Robins, S. P., Shimokomaki, M. & Bailey, A. J.) 771–780
NN′-Diacetylchitobiose, hydrolysis of, by β-*N*-acetylglucosaminidase from *Sclerotinia fructigena* (Reyes, F. & Byrde, R. J. W.) 381–388
1,2-Diacylglycerol, properties and subcellular distribution of a membrane-bound enzyme activity catalysing the breakdown of phosphatidylinositol to D-*myo*-inositol 1:2-cyclic phosphate, D-*myo*-inositol 1-phosphate and, in rat cerebral cortex (Lapetina, E. G. & Michell, R. H.) 433–442
Diamine oxidase, kidney, pig, kinetics of the reaction catalysed by (Bardsley, W. G., Crabbe, M. J. C. & Shindler, J. S.) 459–469
2,5-Diaminoadipic acid lactam residues, inhibition by peptides containing, of DD-carboxypeptidases from *Streptomyces* strains (Nieto, M., Perkins, H. R., Leyh-Bouille, M., Frère, J.-M. & Ghuysen, J.-M.) 163–171
Diet, lipid-rich, effect of, on the properties and composition of lipoprotein particles from the Golgi apparatus of guinea-pig liver (Chapman, M. J., Mills, G. L. & Taylaur, C. E.) 177–185
Diethyl pyrocarbonate, reaction of, with an essential histidine residue in pig heart muscle lactate dehydrogenase (Holbrook, J. J. & Ingram, V. A.) 729–738
p-Dimethylaminomethylbenzylamine, kinetics of the oxidation of, by pig kidney diamine oxidase (Bardsley, W. G., Crabbe, M. J. C. & Shindler, J. S.) 459–469
7,12-Dimethylbenz[*a*]anthracene 5,6-epoxide, chemical synthesis of, and its metabolism by rat liver preparations (Sims, P.) 405–413
2,3-Dimethyl-4-(2-methylenebutyryl)phenoxyacetate, distribution of enzymes that catalyse reactions of reduced glutathione with, and other $\alpha\beta$-unsaturated compounds in the livers of various vertebrate species (Chasseaud, L. F.) 765–769
5,5′-Dithiobis-(2-nitrobenzoate), modification of the activities of acetylcholinesterase and cholinesterase from human blood by, included in the coupled spectrophotometric assay (Brownson, C. & Watts, D. C.) 369–374
Duodenum, mouse, characterization of alternative molecular forms of xanthine oxidase from mouse liver and (Duke, E. J., Joyce, P. & Ryan, J. P.) 187–190

Elastase, pancreas, pig, evidence for the amino acid sequence of (Shotton, D. M. & Hartley, B. S.) 643–675
Embryos, rye-grain, loss of activity of protein-synthesizing systems *in vitro* from, associated with loss of viability of the grains (Roberts, B. E., Payne, P. I. & Osborne, D. J.) 275–286
Enamel, dental, ox, developing, purification and partial characterization of amelogenins from (Eggert, F. M., Allen, G. A. & Burgess, R. C.) 471–484
Endoplasmic reticulum, *see* Reticulum, endoplasmic
Enzymes, shapes of curves of the pH-dependence of the rates of reactions catalysed by (Dixon, H. B. F.) 149–154
Enzymes, significance of abrupt transitions in Lineweaver–Burk plots of the activity of, with particular reference to ox liver glutamate dehydrogenase (Engel, P. C. & Ferdinand, W.) 97–105
Erythrocytes, chicken, turkey and frog, properties of polynucleotides associated with preparations of histones from, and various other sources (Greenaway, P. J. & Murray, K.) 585–591
Erythrocytes, human, modification of the activity of acetylcholinesterase from, by 5,5′-dithiobis-(2-nitrobenzoate) included in the coupled spectrophotometric assay (Brownson, C. & Watts, D. C.) 369–374
Erythrocytes, human, structural requirements for binding to the sugar-transport system of (Barnett, J. E. G., Holman, G. D. & Munday, K. A.) 211–221

Escherichia coli, bacteriophage T4-infected, improved preparation of, for the isolation of polynucleotide kinase (Hughes, S. G. & Brown, P. R.) 583
Escherichia coli (Crookes' strain), purification and characterization of two fructose diphosphate aldolases from (Stribling, D. & Perham, R. N.) 833–841
Escherichia coli K12, competitive inhibition and substrate activity of uridine diphosphate 6-deoxygalactose for uridine diphosphate galactose 4-epimerase from (Spencer, M., Blackburn, P., Ferdinand, W. & Blackburn, G. M.) 421–423
Estradiol, *see* Oestradiol
Estrogen, *see* Oestrogen
Ethanol, characteristics of the 'peroxidatic' reaction of rat liver catalase in the oxidation of (Oshino, N., Oshino, R. & Chance, B.) 555–563
Ethanol, kinetics and mechanism of the reaction catalysed by baker's-yeast alcohol dehydrogenase with, and a variety of other substrates (Dickinson, F. M. & Monger, G. P.) 261–270
Ethylene glycol, effect of neutral salt and, and other alcohols on the thermal stability of soluble and precipitated acid-soluble calf skin collagen (Russell, A. E.) 335–342

Factor X, coagulation, method for the simultaneous isolation of prothrombin and, from ox plasma (Esnouf, M. P., Lloyd, P. H. & Jesty, J.) 781–789
Factor X, coagulation, plasma, ox, preparation of Factor X*a* from, and its action on prothrombin (Jesty, J. & Esnouf, M. P.) 791–797
Factor X*a*, coagulation, ox, determination of the operational molarity of solutions of, by spectrofluorimetric titration (Jameson, G. W., Roberts, D. V., Adams, R. W., Kyle, W. S. A. & Elmore, D. T.) 107–117
Factor X*a*, coagulation, preparation of, from ox plasma Factor X and its action on prothrombin (Jesty, J. & Esnouf, M. P.) 791–797
Fatty acids, long-chain, mechanism of the activation of, catalysed by rat liver microsomal palmitoyl-coenzyme A synthetase (Bar-Tana, J., Rose, G., Brandes, R. & Shapiro, B.) 199–209
Fatty acyl (long-chain)-coenzyme A synthetase, microsomal, liver, rat, structural properties of (Bar-Tana, J. & Rose, G.) 443–449
Ferritin, apo-, spleen, horse, molecular weight of the oligomer of, and its stability to dissociation by dilution (Crichton, R. R., Eason, R., Barclay, A. & Bryce, C. F. A.) 855–857
Ferritin, organ-specificity of, in human and horse liver and spleen (Crichton, R. R., Millar, J. A., Cumming, R. L. C. & Bryce, C. F. A.) 51–59
Fibroblast cells, L929, mouse, cultured, multiple forms of nuclear deoxyribonucleic acid polymerases and their relationship with the cytosol enzyme from (Adams, R. L. P., Henderson, M. A. L., Wood, W. & Lindsay, J. G.) 237–246
Fluorescence, use of, in monitoring the binding of reduced nicotinamide–adenine dinucleotide to lactate dehydrogenases from various muscles (Stinson, R. A. & Holbrook, J. J.) 719–728
Fluorescence, use of, in monitoring the interaction of quinacrine mustard with calf thymus-gland histones (Selander, R.-K. & de la Chapelle, A.) 757–764
Fluorescence, use of, in monitoring the interaction of quinacrine mustard with mononucleotides and polynucleotides (Selander, R.-K.) 749–755
Fluoro-D-glucose 6-phosphates, *see* Deoxyfluoro-D-glucose 6-phosphates
Fructose diphosphate aldolases, two, purification and characterization of, from *Escherichia coli* (Crookes' strain) (Stribling, D. & Perham, R. N.) 833–841
Fructose 1,6-diphosphate, interaction between rabbit liver pyruvate kinase and (Irving, M. G. & Williams, J. F.) 303–313

Gangliosidosis, GM_2-, immunological properties of liver *N*-acetylglucosaminidase from normal humans and from patients with (Carroll, M. & Robinson, D.) 91–96
Genome, coliphage R17, nucleotide sequence containing seventy-three residues from (Rensing, U. F. E.) 593–604
Genome, coliphage R17, nucleotide sequences of similar size from (Rensing, U. F. E., Coulson, A. & Thompson, E. O. P.) 605–610
Germination, loss of activity of protein-synthesizing systems *in vitro* associated with loss of viability of rye grains for (Roberts, B. E., Payne, P. I. & Osborne, D. J.) 275–286
Globin, 3'-hydroxyl-terminal nucleotide sequence of the messenger ribonucleic acid for, from rabbit reticulocytes (Hunt, J. A.) 315–325
Globin, interaction between polyuridylate and the messenger ribonucleic acid for, from rabbit reticulocytes (Hunt, J. A.) 327–333
Glucocerebrosidase, reconstitution of the activity of, from two macromolecular components from human spleen (Ho, M. W., O'Brien, J. S., Radin, N. S. & Erickson, J. S.) 173–176
Glucose 6-phosphate, activation by, of maize-seedling phosphoenolpyruvate carboxylase (Wong, K. F. & Davies, D. D.) 451–458
D-Glucose 6-phosphate, activity of yeast and rat liver glucose 6-phosphate dehydrogenase towards C-2-substituted derivatives of (Bessell, E. M. & Thomas, P.) 83–89
Glucose 6-phosphate dehydrogenase, liver, rat, activity of, towards C-2-substituted derivatives of D-glucose 6-phosphate (Bessell, E. M. & Thomas, P.) 83–89
Glucose 6-phosphate dehydrogenase, yeast, activity of, towards C-2-substituted derivatives of D-glucose 6-phosphate (Bessell, E. M. & Thomas, P.) 83–89
D-Glucose 6-phosphate, effects of C-2-substituted derivatives of, on the activity of yeast glucose phosphate isomerase (Bessell, E. M. & Thomas, P.) 77–82
D-Glucose 6-phosphate–1L-*myo*inositol 1-phosphate cyclase, *see* Inositol cyclase
Glucose phosphate isomerase, yeast, activity of, towards deoxyfluoro-D-glucose 6-phosphates (Bessell, E. M. & Thomas, P.) 77–82
D-Glucose, structural requirements for the binding of, to the sugar-transport system of human erythrocytes (Barnett, J. E. G., Holman, G. D. & Munday, K. A.) 211–221

β-Glucosidase, acid, reconstitution of the activities of glucocerebrosidase and, from two macromolecular components from human spleen (Ho, M. W., O'Brien, J. S., Radin, N. S. & Erickson, J. S.) 173–176

O-Glucuronides, comparison of the properties of the uridine diphosphate glucuronyltransferase activities synthesizing thio-β-D-glucuronides and, in preparations from various animal sources (Illing, H. P. A. & Dutton, G. J.) 139–147

Glutamate dehydrogenase, liver, ox, significance of abrupt transitions in Lineweaver–Burk plots with particular reference to (Engel, P. C. & Ferdinand, W.) 97–105

Glutamate, inhibition by, of maize-seedling phosphoenolpyruvate carboxylase (Wong, K. F. & Davies, D. D.) 451–458

Glutathione (reduced) *S*-alkenetransferases, distribution of, in the livers of various vertebrate species (Chasseaud, L. F.) 765–769

Glutathione, reduced, distribution of enzymes that catalyse reactions of, with αβ-unsaturated compounds in the livers of various vertebrate species (Chasseaud, L. F.) 765–769

Glycerol, effect of neutral salt and, and other alcohols on the thermal stability of soluble and precipitated acid-soluble calf skin collagen (Russell, A. E.) 335–342

Glycine max, *see* Soya bean

Glycopeptides, isolation of a new neuraminic acid derivative and three types of, from the Cuvierian tubules of the sea-cucumber *Holothuria forskali* (Isemura, M., Zahn, R. K. & Schmid, K.) 509–521

Glycosaminoglycans, differences in, of proteolgycans extracted from ox nasal cartilage with solutions of high and of low ionic strengths (Mayes, R. W., Mason, R. M. & Griffin, D. C.) 541–553

Glycosaminoglycans, quantitative determination of, in urine with Alcian Blue 8GX (Whiteman, P.) 351–357

Glycosaminoglycans, quantitative measurement of complexes of Alcian Blue 8GX with (Whiteman, P.) 343–350

Golgi apparatus, liver, guinea-pig, effect of a lipid-rich diet on the properties and composition of lipoprotein particles from (Chapman, M. J., Mills, G. L. & Taylaur, C. E.) 177–185

p-Guanidinobenzoic acid, determination of the operational molarity of solutions of serine proteinases by spectrofluorimetric titration with the 4-methylumbelliferyl ester of (Jameson, G. W., Roberts, D. V., Adams, R. W., Kyle, W. S. A. & Elmore, D. T.) 107–117

Guanine deaminase, liver and brain, rat and mouse, regulation of the activities of the isoenzymes of (Sree Kumar, K., Sitaramayya, A. & Krishnan, P. S). 683–687

Guanosine triphosphate, effects of, on the activities of rat and mouse liver and brain guanine deaminase activities (Sree Kumar, K., Sitaramayya, A. & Krishnan, P. S.) 683–687

Haemoglobin, 3′-hydroxyl-terminal nucleotide sequence of the messenger ribonucleic acid for, from rabbit reticulocytes (Hunt, J. A.) 315–325

Haemoglobin, interaction between polyuridylate and the messenger ribonucleic acid for, from rabbit reticulocytes (Hunt, J. A.) 327–333

Haptoglobin, human, lack of effect of, on the activity of human liver cathepsin B1 (Starkey, P. M. & Barrett, A. J.) 823–831

Heart, horse, accurate determination of the molecular weight of cytochrome *c* from, with a sensitive polyacrylamide-gel osmometer (Ogston, A. G. & Preston, B. N.) 843–850

Heart, horse, effect of steady illumination on the binding of carbon monoxide by carboxymethylated cytochrome *c* from (Wilson, M. T., Brunori, M., Bonaventura, J. & Bonaventura, C.) 863–865

Heart muscle, pig, equilibrium binding of nicotinamide–adenine dinucleotide to lactate dehydrogenases from, and various other muscles (Stinson, R. A. & Holbrook, J. J.) 719–728

Heart muscle, pig, ionic properties of an essential histidine residue in lactate dehydrogenase from (Holbrook, J. J. & Ingram, V. A.) 729–738

Heart muscle, pig, use of ternary complexes to study ionizations and isomerizations during catalysis by lactate dehydrogenase from (Holbrook, J. J. & Stinson, R. A.) 739–748

Heavy meromyosin, *see* Meromyosin, heavy

Hepatoma cell, ascites, rat, characterization of a non-histone protein isolated from histone F1 preparations from, and other sources (Smith, J. A. & Stocken, L. A.) 859–861

Hexosaminidase, *see* *N*-Acetyl-β-D-glucosaminidase

D-*xylo*-5-Hexosulose 6-phosphate, evidence for the formation of, as an enzyme-bound intermediate in the conversion of D-glucose 6-phosphate into 1L-*myo*inositol 1-phosphate by rat testis inositol cyclase (Barnett, J. E. G., Rasheed, A. & Corina, D. L.) 21–30

Histidine residue, essential, ionic properties of, in pig heart muscle lactate dehydrogenase (Holbrook, J. J. & Ingram, V. A.) 729–738

Histone F1, characterization of a non-histone protein isolated from preparations of, from rat liver, thymus gland and ascites hepatoma cells (Smith, J. A. & Stocken, L. A.) 859–861

Histones, properties of polynucleotides associated with preparations of, from various sources (Greenaway, P. J. & Murray, K.) 585–591

Histones, thymus-gland, calf, interaction of quinacrine mustard with (Selander, R.-K. & de la Chapelle, A.) 757–764

Holothuria forskali, *see* Sea-cucumber

Hurler's syndrome, quantitative determination with Alcian Blue 8GX of glycosaminoglycans in the urine of human patients with (Whiteman, P.) 351–357

Hydrogen ions, uptake of, during the reaction catalysed by pig heart-muscle and skeletal-muscle lactate dehydrogenases (Holbrook, J. J. & Stinson, R. A.) 739–748

Hydrogen isotope (^2H), effect of the presence of, on the kinetics of the oxidation of *p*-dimethylaminomethyl[αα-$^2H_2$]benzylamine by pig kidney diamine oxidase (Bardsley, W. G., Crabbe, M. J. C. & Shindler, J. S.) 459–469

Hydrogen peroxide, generation of, during the oxidation of ethanol by the 'peroxidatic' activity of rat liver catalase (Oshino, N., Oshino, R. & Chance, B.) 555–563
Hydroxylysine residues, identification of, as the site of attachment of carbohydrate residues in glycopeptides isolated from the Cuvierian tubules of the sea-cucumber *Holothuria forskali* (Isemura, M., Zahn, R. K. & Schmid, K.) 509–521
7-Hydroxymethyl-12-methylbenz[*a*]anthracene 5,6-epoxide, chemical synthesis of, and its metabolism by rat liver preparations (Sims, P.) 405–413

Immunoglobulin G, human, inhibition by, of the activity of human liver cathepsin B1 (Starkey, P. M. & Barrett, A. J.) 823–831
Immunoglobulin M, human, investigations of the structural function of the J chain in (Ricardo, M. J., Jr. & Inman, F. P.) 677–682
Inositol cyclase, testis, rat, partial reactions of (Barnett, J. E. G., Rasheed, A. & Corina, D. L.) 21–30
D-*myo*Inositol 1:2-cyclic phosphate, properties and subcellular distribution of a membrane-bound enzyme activity catalysing the breakdown of phosphatidylinositol to 1,2-diacylglycerol, D-*myo*inositol 1-phosphate and, in rat cerebral cortex (Lapetina, E. G. & Michell, R. H.) 433–442
D-*myo*Inositol 1-phosphate, properties and subcellular distribution of a membrane-bound enzyme activity catalysing the breakdown of phosphatidylinositol to 1,2-diacylglycerol, D-*myo*inositol 1:2-cyclic phosphate and, in rat cerebral cortex (Lapetina, E. G. & Michell, R. H.) 433–442
Insulin, ox, reaction of methyl iodopicolinimidate with the amino groups of (Riley, M. & Perham, R. N.) 625–635
Intestine, small, mouse, characterization of alternative molecular forms of xanthine oxidase from mouse liver and (Duke, E. J., Joyce, P. & Ryan, J. P.) 187–190
Intestine, small, rat, partial purification and properties of tryptophan 5-hydroxylase from (Noguchi, T., Nishino, M. & Kido, R.) 375–380
Ionic strength, effects of dielectric constant and, on the kinetics of the oxidation of *p*-dimethylaminomethyl[$\alpha\alpha$-$^2H_2$]benzylamine by pig kidney diamine oxidase (Bardsley, W. G., Crabbe, M. J. C. & Shindler, J. S.) 459–469
Ionic strengths, high and low, composition of proteoglycans extracted from ox nasal cartilage with solutions of (Mayes, R. W., Mason, R. M. & Griffin, D. C.) 541–553
Ionic strengths, various, extraction of proteoglycans from ox nasal cartilage with inorganic salt solutions of (Mason, R. M. & Mayes, R. W.) 535–540
Isocoproporphyrin, determination of the structure of, with the aid of nuclear-magnetic-resonance- and mass-spectral methods (Stoll, M. S., Elder, G. H., Games, D. E., O'Hanlon, P., Millington, D. S. & Jackson, A. H.) 429–432

'Joining' chain (J chain), investigations of the structural function of, in human immunoglobulin M (Ricardo, M. J., Jr. & Inman, F. P.) 677–682

Keratan sulphate, extraction of proteoglycan enriched in, from ox nasal cartilage with solutions of high ionic strength (Mayes, R. W., Mason, R. M. & Griffin, D. C.) 541–553
5-Keto-D-glucose 6-phosphate, *see* D-*xylo*-5-Hexosulose 6-phosphate
Kidney cells, baby-hamster (BHK-21/C13), biosynthesis of deoxyribonucleic acid in (Hayton, G. J., Pearson, C. K., Scaife, J. R. & Keir, H. M.) 499–508
Kidney, mouse, properties of the uridine diphosphate glucuronyltransferase activity synthesizing thio-β-D-glucuronides in preparations of (Illing, H. P. A. & Dutton, G. J.) 139–147
Kidney, ox, comparison of the kinetic properties of partially purified pyruvate dehydrogenase complex from ox cerebral cortex and (Blass, J. P. & Lewis, C. A.) 31–37
Kidney, ox, inhibition by acetaldehyde of the pyruvate dehydrogenase complex from (Blass, J. P. & Lewis C. A.) 415–416
Kidney, pig, inhibition by oxidized or reduced nicotinamide–adenine dinucleotide and related compounds of alkaline phosphatase from (Ramasamy, I. & Butterworth, P. J.) 359–367
Kidney, pig, kinetics of the reaction catalysed by diamine oxidase from (Bardsley, W. G., Crabbe, M. J. C. & Shindler, J. S.) 459–469

Lactate dehydrogenase, muscle, heart, pig, ionic properties of an essential histidine residue in (Holbrook, J. J. & Ingram, V. A.) 729–738
Lactate dehydrogenases, muscle, heart and skeletal, pig, use of ternary complexes to study ionizations and isomerizations during catalysis by (Holbrook, J. J. & Stinson, R. A.) 739–748
Lactate dehydrogenases, muscle, various, equilibrium binding of nicotinamide–adenine dinucleotide to (Stinson, R. A. & Holbrook, J. J.) 719–728
Lactate, purification and characterization of two fructose diphosphate aldolases from *Escherichia coli* (Crookes' strain) grown on pyruvate or (Stribling, D. & Perham, R. N.) 833–841
Lanthanum chloride, extraction of proteoglycans from ox nasal cartilage with solutions containing (Mason, R. M. & Mayes, R. W.) 535–540
Lectins, wheat-germ, purification, composition and specificity of agglutinins from (Allen, A. K., Neuberger, A. & Sharon, N.) 155–162
Leek (*Allium porrum*), amino acid sequence of cytochrome *c* from (Brown, R. H. & Boulter, D.) 247–251
Lipid, effect of a diet rich in, on the properties and composition of lipoprotein particles from the Golgi apparatus of guinea-pig liver (Chapman, M. J., Mills, G. L. & Taylaur, C. E.) 177–185
Lipoprotein particles, effect of a lipid-rich diet on the properties and composition of, from the Golgi apparatus of guinea-pig liver (Chapman, M. J., Mills, G. L. & Taylaur, C. E.) 177–185
Lipoproteins, very-low-density, serum, guinea-pig, evidence that lipoprotein particles of the Golgi apparatus of the liver are precursors of (Chapman, M. J., Mills, G. L. & Taylaur, C. E.) 177–185

Liver, distribution of enzymes that catalyse reactions of reduced glutathione with $\alpha\beta$-unsaturated compounds in, of various vertebrate species (Chasseaud, L. F.) 765–769
Liver, guinea-pig, effect of a lipid-rich diet on the properties and composition of lipoprotein particles from the Golgi apparatus of (Chapman, M. J., Mills, G. L. & Taylaur, C. E.) 177–185
Liver, human and horse, organ-specificity of ferritin in, and spleen (Crichton, R. R., Millar, J. A., Cumming, R. L. C. & Bryce, C. F. A.) 51–59
Liver, human, immunological properties of *N*-acetyl-β-D-glucosaminidase from, of normal subjects and of patients with GM_2-gangliosidosis (Carroll, M. & Robinson, D.) 91–96
Liver, human, inhibition by α_2-macroglobulin and other human serum proteins of the activity of cathepsin B1 from (Starkey, P. M. & Barrett, A. J.) 823–831
Liver, human, purification and properties of cathepsin B1 from (Barrett, A. J.) 809–822
Liver, mouse, characterization of alternative molecular forms of xanthine oxidase from mouse duodenum and (Duke, E. J., Joyce, P. & Ryan, J. P.) 187–190
Liver, mouse, rat and chick-embryo, properties of the uridine diphosphate glucuronyltransferase activity synthesizing thio-β-D-glucuronides in preparations of (Illing, H. P. A. & Dutton, G. J.) 139–147
Liver, ox, significance of abrupt transitions in Lineweaver–Burk plots with particular reference to glutamate dehydrogenase from (Engel, P. C. & Ferdinand, W.) 97–105
Liver, rabbit, interaction between pyruvate kinase from, and its allosteric effector fructose 1,6-diphosphate (Irving, M. G. & Williams, J. F.) 303–313
Liver, rabbit, kinetic studies on the regulation of pyruvate kinase from (Irving, M. G. & Williams, J. F.) 287–301
Liver, rat, activity of glucose 6-phosphate dehydrogenase from, towards C-2-substituted derivatives of D-glucose 6-phosphate (Bessell, E. M. & Thomas, P.) 83–89
Liver, rat and mouse, regulation of the activities of guanine deaminase isoenzymes in rat and mouse brain and (Sree Kumar, K., Sitaramayya, A. & Krishnan, P. S.) 683–687
Liver, rat, binding of polyribosomes to smooth and rough endoplasmic-reticulum membranes from (Rolleston, F. S. & Mark, D.) 851–853
Liver, rat, characteristics of the 'peroxidatic' reaction of catalase from, in the oxidation of ethanol (Oshino, N., Oshino, R. & Chance, B.) 555–563
Liver, rat, characterization of a non-histone protein isolated from histone F1 preparations from, and other sources (Smith, J. A. & Stocken, L. A.) 859–861
Liver, rat, mechanism of the activation of long-chain fatty acids catalysed by microsomal palmitoyl-coenzyme A synthetase from (Bar-Tana, J., Rose, G., Brandes, R. & Shapiro, B.) 199–209
Liver, rat, metabolism of epoxides related to 7,12-dimethylbenz[*a*]anthracene by preparations of (Sims, P.) 405–413
Liver, rat, stimulation by ascorbate of the activity of phenylalanine hydroxylase from (Stone, K. J. & Townsley, B. H.) 611–613
Liver, rat, structural properties of microsomal palmitoyl-coenzyme A synthetase from (Bar-Tana, J. & Rose, G.) 443–449
Lysine residues, additional, reactivity towards methyl picolinimidate of, in the coat protein of mutants of tobacco-mosaic virus and the reagent's usefulness as a probe for investigating the topography of the virus (Perham, R. N.) 119–126
Lysosomes, blood-platelet, human, localization of phospholipase A_1 in (Smith, J. B., Silver, M. J. & Webster, G. R.) 615–618

Macroglobulin, *see* Immunoglobulin M
α_2-Macroglobulin, serum, human, inhibition by, of the activity of human liver cathepsin B1 (Starkey, P. M. & Barrett, A. J.) 823–831
Magnesium ions, effects of, on the activities of rat and mouse liver and brain guanine deaminase isoenzymes (Sree Kumar, K., Sitaramayya, A. & Krishnan, P. S.) 683–687
Maize (*Zea mays*) seedlings, regulation by metabolites of the activity of phosphoenolpyruvate carboxylase from (Wong, K. F. & Davies, D. D.) 451–458
Malayan pit viper (*Agkistrodon rhodostoma*), isolation and properties of the coagulant enzyme Arvin from the venom of (Hatton, M. W. C.) 799–807
Membranes, endoplasmic-reticulum, liver, rat, smooth and rough, binding of polyribosomes to (Rolleston, F. S. & Mak, D.) 851–853
Membranes, plasma, properties of an enzyme activity bound to, catalysing the breakdown of phosphatidylinositol to 1,2-diacylglycerol, D-*myo*inositol 1:2-cyclic phosphate and D-*myo*inositol 1-phosphate in rat cerebral cortex (Lapetina, E. G. & Michell, R. H.) 433–442
2-Mercaptoethylamine, dissociation of human immunoglobulin M by treatment with (Ricardo, M. J., Jr. & Inman, F. P.) 677–682
Meromyosin, heavy, skeletal-muscle, rabbit, isolation and properties of subfragment 1 preparations obtained by proteolysis with papain from (Stone, D. & Perry, S. V.) 127–137
Messenger ribonucleic acid, *see* Ribonucleic acid, messenger
Methyl iodopicolinimidate, reaction of, with protein amino groups and its possible use as a general method for preparing isomorphous heavy-atom derivatives of proteins (Riley, M. & Perham, R. N) 625–635
Methyl 5-iodopyridine-2-carboximidate, reaction of, with protein amino groups and its possible use as a general method for preparing isomorphous heavy-atom derivatives of proteins (Riley, M. & Perham, R. N.) 625–635
Methyl picolinimidate, reactivity towards, of additional lysine residues in the coat protein of mutants of tobacco-mosaic virus and its usefulness as a probe for investigating the topography of the virus (Perham, R. N.) 119–126
Methyl vinyl sulphone, distribution of enzymes that catalyse reactions of reduced glutathione with, and other $\alpha\beta$-unsaturated compounds in the livers of various vertebrate species (Chasseaud, L. F.) 765–769

2′-*O*-Methyladenylyl-2′-*O*-methyluridylylguanylate, preparation of an alkali-stable trinucleotide from yeast ribonucleic acid and its identification as (Trim, A. R. & Parker, J. E.) 257–259
4-Methylumbelliferone, determination of the operational molarity of solutions of serine proteinases by spectrofluorimetric titration with esters of (Jameson, G. W., Roberts, D. V., Adams, R. W., Kyle, W. S. A. & Elmore, D. T.) 107–117
Micrococcus denitrificans, purification and properties of 5-aminolaevulinate synthetase from (Tait, G. H.) 389–403
Micrococcus lysodeikticus N.C.T.C. 2665, selective extraction of teichuronic acids from the cell walls of, and other Gram-positive bacteria (Pavlik, J. G. & Rogers, H. J.) 619–621
Microsomal fraction, liver, rat, mechanism of the activation of long-chain fatty acids catalysed by palmitoyl-coenzyme A synthetase of (Bar-Tana, J., Rose, G., Brandes, R. & Shapiro, B.) 199–209
Microsomal fraction, liver, rat, metabolism of epoxides related to 7,12-dimethylbenz[*a*]anthracene by (Sims, P.) 405–413
Microsomal fraction, liver, rat, structural properties of palmitoyl-coenzyme A synthetase from (Bar-Tana, J. & Rose, G.) 443–449
Microsomal preparations, properties of the uridine diphosphate glucuronyltransferase activity synthesizing thio-β-D-glucuronides in, from various animal sources (Illing, H. P. A. & Dutton, G. J.) 139–147
Milk, cow, purification, interconversion and properties of the type D (dehydrogenase) and type O (oxidase) forms of xanthine oxidase from (Battelli, M. G., Lorenzoni, E. & Stirpe, F.) 191–198
Mucopolysaccharidoses, quantitative determination with Alcian Blue 8GX of glycosaminoglycans in the urine of human patients with (Whiteman, P.) 351–357
Muscle, heart, pig, equilibrium binding of nicotinamide–adenine dinucleotide to lactate dehydrogenases from, and various other muscles (Stinson, R. A. & Holbrook, J. J.) 719–728
Muscle, heart, pig, ionic properties of an essential histidine residue in lactate dehydrogenase from (Holbrook, J. J. & Ingram, V. A.) 729–738
Muscle, heart, pig, use of ternary complexes to study ionizations and isomerizations during catalysis by lactate dehydrogenase from (Holbrook, J. J. & Stinson, R. A.) 739–748
Muscle, skeletal, pig, use of ternary complexes to study ionizations and isomerizations during catalysis by lactate dehydrogenase from (Holbrook, J. J. & Stinson, R. A.) 739–748
Muscle, skeletal, rabbit, isolation and properties of subfragment 1 preparations obtained by proteolysis with papain from the heavy chain of myosin from (Stone, D. & Perry, S. V.) 127–137
Muscle, skeletal, rabbit, kinetics of the reaction catalysed by pyruvate kinase from (Ainsworth, S. & Macfarlane, N.) 223–236
Muscle, skeletal, rabbit, phosphorylation of the '37000 component' (troponin-T) of the troponin complex from (Perry, S. V. & Cole, H. A.) 425–428
Muscle, skeletal, vertebrate, equilibrium binding of nicotinamide–adenine dinucleotide to lactate dehydrogenases from, and various other muscles (Stinson, R. A. & Holbrook, J. J.) 719–728
Muscle, tail, lobster, equilibrium binding of nicotinamide–adenine dinucleotide to lactate dehydrogenases from, and various other muscles (Stinson, R. A. & Holbrook, J. J.) 719–728
Mycobacillin, stereoconfiguration of the amino acid residues in peptides isolated from a *Bacillus subtilis* B_3 cell-free system synthesizing (Sengupta, S. & Bose, S. K.) 623–624
Myosin, skeletal-muscle, rabbit, isolation and properties of subfragment 1 preparations obtained by proteolysis with papain from the heavy chain of (Stone, D. & Perry, S. V.) 127–137

Neuraminic acid, isolation of three types of glycopeptides and a new derivative of, from the Cuvierian tubules of the sea-cucumber *Holothuria forskali* (Isemura, M., Zahn, R. K. & Schmid, K.) 509–521
Nicotinamide, inhibition by, of pig kidney alkaline phosphatase (Ramasamy, I. & Butterworth, P. J.) 359–367
Nicotinamide–adenine dinucleotide, oxidized and reduced, inhibition by, of pig kidney alkaline phosphatase (Ramasamy, I. & Butterworth, P. J.) 359–367
Nicotinamide–adenine dinucleotide, reduced and oxidized, equilibrium binding of, to lactate dehydrogenases from various muscles (Stinson, R. A. & Holbrook, J. J.) 719–728
Nitrogen, kinetics of the reduction of acetylene and, by the nitrogenase complex from soya-bean root-nodule bacteroids (Bergersen, F. J. & Turner, G. L.) 61–75
Nitrogenase complex, kinetic properties of, from soya-bean root-nodule bacteroids (Bergersen, F. J. & Turner, G. L.) 61–75
o-Nitrophenylpyruvate, ionizations and isomerizations during the reduction of, catalysed by pig heart-muscle and skeletal-muscle lactate dehydrogenases (Holbrook, J. J. & Stinson, R. A.) 739–748
3-(β-Nitrovinyl)indole, distribution of enzymes that catalyse reactions of reduced glutathione with, and other αβ-unsaturated compounds in the livers of various vertebrate species (Chasseaud, L. F.) 765–769
Nuclear ribonucleic acid, *see* Ribonucleic acid, nuclear
Nuclei, multiple forms of deoxyribonucleic acid polymerases from, and their relationship with the cytosol enzyme from cultured mouse L929 cells (Adams, R. L. P., Henderson, M. A. L., Wood, W. & Lindsay, J. G.) 237–246
Nucleotide sequence containing seventy-three residues from the coliphage R17 genome (Rensing, U. F. E.) 593–604
Nucleotide sequence, 3′-hydroxyl-terminal, of globin messenger ribonucleic acid from rabbit reticulocytes (Hunt, J. A.) 315–325
Nucleotide sequences, 5′-hydroxyl-terminal, use of polynucleotide kinase and nucleotide 'mapping' methods for the determination of, of deoxyribonucleic acids from bacteriophages λ and 424 (Murray, K.) 569–582

Nucleotide sequences of similar size from the coliphage R17 genome (Rensing, U. F. E., Coulson, A. & Thompson, E. O. P.) 605–610
Nucleotides, interaction of quinacrine mustard with (Selander, R.-K.) 749–755

Oestradiol-17β, stimulation by, of the biosynthesis of heterogeneous nuclear ribonucleic acid in immature rat uterus (Knowler, J. T. & Smellie, R. M. S.) 689–697
Oestrogen, stimulation by, of the biosynthesis of heterogeneous nuclear ribonucleic acid in immature rat uterus (Knowler, J. T. & Smellie, R. M. S.) 689–697
Osmometer, polyacrylamide-gel, sensitive, accurate determination of the molecular weights of proteins and other macromolecules with (Ogston, A. G. & Preston, B. N.) 843–850
α-Oxo acids, effects of, on partially purified ox cerebral-cortex pyruvate dehydrogenase complex (Blass, J. P. & Lewis, C. A.) 31–37
2-Oxoglutarate, inhibition by, of maize-seedling phosphoenolpyruvate carboxylase (Wong, K. F. & Davies, D. D.) 451–458

Palmitoyl-coenzyme A synthetase, microsomal, liver, rat, mechanism of the activation of long-chain fatty acids catalysed by (Bar-Tana, J., Rose, G., Brandes, R. & Shapiro, B.) 199–209
Palmitoyl-coenzyme A synthetase, microsomal, liver, rat, structural properties of (Bar-Tana, J. & Rose, G.) 443–449
Pancreas, pig, evidence for the amino acid sequence of elastase from (Shotton, D. M. & Hartley, B. S.) 643–675
Pea (*Pisum sativum*) seedlings, nature of the polydisperse ribonucleic acid obtained from (Jackson, M. & Ingle, J.) 523–533
Peptides, stereoconfiguration of the amino acid residues in, isolated from a mycobacillin-synthesizing cell-free system prepared from *Bacillus subtilis* B_3 (Sengupta, S. & Bose, S. K.) 623–624
Peptides, synthesis of, that inhibit DD-carboxypeptidases from *Streptomyces* strains (Nieto, M., Perkins, H. R., Leyh-Bouille, M., Frère, J.-M. & Ghuysen, J.-M.) 163–171
Peptidyltransferases, specificity of *Streptomyces* DD-carboxypeptidases acting as, for amino compounds as carboxyl acceptors (Perkins, H. R., Nieto, M., Frère, J.-M., Leyh-Bouille, M. & Ghuysen, J.-M.) 707–718
'Peroxidatic' reaction, characteristics of, of rat liver catalase in the oxidation of ethanol (Oshino, N., Oshino, R. & Chance, B.) 555–563
pH, shapes of curves of the dependence of reaction rates on (Dixon, H. B. F.) 149–154
Phage, *see* Bacteriophage; *see also* Coliphage
Phenylalanine hydroxylase, liver, rat, stimulation by ascorbate of the activity of (Stone, K. J. & Townsley, B. H.) 611–613
Phosphatase, alkaline, kidney, pig, inhibition of, by oxidized and reduced nicotinamide–adenine dinucleotide and related compounds (Ramasamy, I. & Butterworth, P. J.) 359–367
Phosphatidylinositol, properties and subcellular distribution of a membrane-bound enzyme activity catalysing the breakdown of, to 1,2-diacylglycerol, D-*myo*inositol 1:2-cyclic phosphate and D-*myo*inositol 1-phosphate in rat cerebral cortex (Lapetina, E. G. & Michell, R. H.) 433–442
Phosphoenolpyruvate carboxylase, maize-seedling, regulation by metabolites of the activity of (Wong, K. F. & Davies, D. D.) 451–458
Phosphoenolpyruvate, kinetics of the reaction between adenosine diphosphate and, catalysed by rabbit skeletal-muscle pyruvate kinase (Ainsworth, S. & Macfarlane, N.) 223–236
Phospholipase A_1, activity of, in human blood platelets (Smith, J. B., Silver, M. J. & Webster, G. R.) 615–618
Phosphorylase *b* kinase, phosphorylation of the '37000 component' (troponin-T) from rabbit skeletal-muscle troponin complex by (Perry, S. V. & Cole, H. A.) 425–428
Phycocyanin, C-, characterization of, from the extremely halo-tolerant blue–green alga *Coccochloris elabens* (Kao, O. H. W., Berns, D. S. & Town, W. R.) 39–50
Pisum sativum, *see* Pea
Plasma albumin, *see* Albumin, plasma
Plasma, human, modification of the activity of cholinesterase from, by 5,5′-dithiobis-(2-nitrobenzoate) included in the coupled spectrophotometric assay (Brownson, C. & Watts, D. C.) 369–374
Plasma membrane, *see* Membrane, plasma
Plasma, ox, method for the simultaneous isolation of Factor X and prothrombin from (Esnouf, M. P., Lloyd, P. H. & Jesty, J.) 781–789
Plasma, ox, preparation of Factor X*a* from Factor X and its action on prothrombin from (Jesty, J. & Esnouf, M. P.) 791–797
Platelets, blood, human, activity of phospholipase A_1 in (Smith, J. B., Silver, M. J. & Webster, G. R.) 615–618
Policy of the Journal and instructions to authors 1–20
Polyadenylate sequences, interaction between polyuridylate and, of globin messenger ribonucleic acid from rabbit reticulocytes (Hunt, J. A.) 327–333
Polyadenylate sequences, presence of, at the 3′-hydroxyl terminus of globin messenger ribonucleic acid from rabbit reticulocytes (Hunt, J. A.) 315–325
Polynucleotide kinase, improved preparation of bacteriophage T4-infected *Escherichia coli* for the isolation of (Hughes, S. G. & Brown, P. R.) 583
Polynucleotide kinase, use of, and nucleotide 'mapping' methods for the determination of the 5′-hydroxy-terminal nucleotide sequences of deoxyribonucleic acids from bacteriophages λ and 424 (Murray, K.) 569–582
Polynucleotides, interaction of quinacrine mustard with (Selander, R.-K.) 749–755
Polynucleotides, properties of, associated with preparations of histones from various sources (Greenaway, P. J. & Murray, K.) 585–591
Polyribosomes, binding of, to smooth and rough rat liver endoplasmic-reticulum membranes (Rolleston, F. S. & Mak, D.) 851–853
Polysomes, *see* Polyribosomes

Polyuridylate, interaction between, and globin messenger ribonucleic acid from rabbit reticulocytes (Hunt, J. A.) 327–333
Porphyrins, use of nuclear-magnetic-resonance- and mass-spectral methods for the determination of the structure of (Stoll, M. S., Elder, G. H., Games, D. E., O'Hanlon, P., Millington, D. S. & Jackson, A. H.) 429–432
Potassium chloride, neutral, effect of alcohols and, on the thermal stability of soluble and precipitated acid-soluble calf skin collagen (Russell, A. E.) 335–342
Propan-1-ol, kinetics and mechanism of the reaction catalysed by baker's-yeast alcohol dehydrogenase with, and a variety of other substrates (Dickinson, F. M. & Monger, G. P.) 261–270
Propan-2-ol, kinetics and mechanism of the reaction catalysed by baker's-yeast alcohol dehydrogenase with, and a variety of other substrates (Dickinson, F. M. & Monger, G. P.) 261–270
Protamine, interaction of quinacrine mustard with, and other proteins (Selander, R.-K. & de la Chapelle, A.) 757–764
Proteases, *see* Proteinases
Protein, coat, reactivity towards methyl picolinimidate of additional lysine residues in, of mutants of tobacco-mosaic virus and the reagent's usefulness as a probe for investigating the topography of the virus (Perham, R. N.) 119–126
Protein kinase, activity of, in commercially available crystalline preparations of yeast alcohol dehydrogenase (Kemp, B. E., Froscio, M. & Murray, A. W.) 271–274
Protein, loss of activity of systems synthesizing, *in vitro* associated with loss of viability of rye grains (Roberts, B. E., Payne, P. I. & Osborne, D. J.) 275–286
Protein, non-histone, characterization of, isolated from histone F1 preparations from rat liver, thymus gland and ascites-hepatoma cells (Smith, J. A. & Stocken L. A.) 859–861
Proteinases, serine, ox, determination of the operational molarity of solutions of, by spectrofluorimetric titration (Jameson, G. W., Roberts, D. V., Adams, R. W., Kyle, W. S. A. & Elmore, D. T.) 107–117
Protein–polysaccharide, *see* Proteoglycan
Proteins, accurate determination of the molecular weights of, with a sensitive polyacrylamide-gel osmometer (Ogston, A. G. & Preston, B. N.) 843–850
Proteins, interaction of quinacrine mustard with (Selander, R.-K. & de la Chapelle, A.) 757–764
Proteins, possible use of the reaction of methyl iodopicolinimidate with protein amino groups as a general method for preparing isomorphous heavy-atom derivatives of (Riley, M. & Perham, R. N.) 625–635
Proteins, purification and partial characterization of, from developing ox dental enamel (Eggert, F. M., Allen, G. A. & Burgess, R. C.) 471–484
Proteins, serum, human, inhibition by, of the activity of human liver cathepsin B1 (Starkey, P. M. & Barrett, A. J.) 823–831
Proteins, use of 3-bromopropionic acid for the determination of thiol groups of (Bradbury, A. F. & Smyth, D. G.) 637–642
Proteoglycans, composition of, extracted from ox nasal cartilage with solutions of high and of low ionic strengths (Mayes, R. W., Mason, R. M. & Griffin, D. C.) 541–553
Proteoglycans, extraction of, from ox nasal cartilage with inorganic salt solutions of various ionic strengths (Mason, R. M. & Mayes, R. W.) 535–540
Proteolysis, conversion of the mouse liver form of xanthine oxidase into the duodenal form by (Duke, E. J., Joyce, P. & Ryan, J. P.) 187–190
Proteolysis, irreversible formation of a type O (oxidase) form from cow's-milk xanthine oxidase by (Battelli, M. G., Lorenzoni, E. & Stripe, F.) 191–198
Prothrombin, method for the simultaneous isolation of Factor X and, from ox plasma (Esnouf, M. P., Lloyd, P. H. & Jesty, J.) 781–789
Prothrombin, plasma, ox, preparation of Factor X*a* from Factor X and its action on (Jesty, J. & Esnouf, M. P.) 791–797
Protonation, effect of, of reactants or intermediates on the pH-dependence of reaction rates (Dixon, H. B. F.) 149–154
Protons, uptake of, during the reaction catalysed by pig heart muscle and skeletal-muscle lactate dehydrogenases (Holbrook, J. J. & Stinson, R. A.) 739–748
Pseudocholinesterase, *see* Cholinesterase
Pseudomonas fluorescens A.T.C.C. 17400, amino acid sequence of cytochrome *c*-551 from (Ambler, R. P. & Wynn, M.) 485–498
Pseudomonas mendocina, amino acid sequence of cytochrome *c*-551 from (Ambler, R. P. & Wynn, M.) 485–498
Pseudomonas stutzeri A.T.C.C. 17588, amino acid sequence of cytochrome *c*-551 from (Ambler, R. P. & Wynn, M.) 485–498
Pyruvate dehydrogenase complex, cerebral-cortex and kidney, ox, inhibition by acetaldehyde of (Blass, J. P. & Lewis, C. A.) 415–416
Pyruvate dehydrogenase complex, cerebral-cortex, ox, partially purified, kinetic properties of (Blass, J. P. & Lewis, C. A.) 31–37
Pyruvate kinase, liver, rabbit, interaction between, and its allosteric effector fructose 1,6-diphosphate (Irving, M. G. & Williams, J. F.) 303–313
Pyruvate kinase, liver, rabbit, kinetic studies on the regulation of (Irving, M. G. & Williams, J. F.) 287–301
Pyruvate kinase, skeletal-muscle, rabbit, kinetics of the reaction catalysed by (Ainsworth, S. & Macfarlane, N.) 223–236
Pyruvate, purification and characterization of two fructose diphosphate aldolases from *Escherichia coli* (Crookes' strain) grown on lactate or (Stribling, D. & Perham, R. N.) 833–841

Quinacrine mustard, interaction of, with calf thymus-gland histones (Selander, R.-K. & de la Chapelle, A.) 757–764
Quinacrine mustard, interaction of, with mononucleotides and polynucleotides (Selander, R.-K.) 749–755

Red blood cells, *see* Erythrocytes
Reticulocytes, rabbit, 3′-hydroxyl-terminal nucleotide sequence of globin messenger ribonucleic acid from (Hunt, J. A.) 315–325
Reticulocytes, rabbit, interaction between polyuridylate and globin messenger ribonucleic acid from (Hunt, J. A.) 327–333
Reticulum, endoplasmic, liver, rat, smooth and rough, binding of polyribosomes to membranes of (Rolleston F. S. & Mak, D.) 851–853
Ribonucleic acid, genome, coliphage R17, nucleotide sequence containing seventy-three residues from (Rensing, U. F. E.) 593–604
Ribonucleic acid, genome, coliphage R17, nucleotide sequences of similar size from (Rensing, U. F. E., Coulson, A. & Thompson, E. O. P.) 605–610
Ribonucleic acid, messenger, globin, 3′-hydroxyl-terminal nucleotide sequence of, from rabbit reticulocytes (Hunt, J. A.) 315–325
Ribonucleic acid, messenger, globin, interaction between polyuridylate and, from rabbit reticulocytes (Hunt, J. A.) 327–333
Ribonucleic acid, nuclear, heterogeneous, oestrogen-stimulated biosynthesis of, in immature rat uterus (Knowler, J. T. & Smellie, R. M. S.) 689–697
Ribonucleic acid, polydisperse, nature of, obtained from pea seedlings (Jackson, M. & Ingle, J.) 523–533
Ribonucleic acid, ribosomal, pea-seedling, effect of pH on the aggregation of, during chromatography on methylated albumin on kieselguhr (Jackson, M. & Ingle, J.) 523–533
Ribonucleic acid, yeast, preparation of an alkali-stable trinucleotide from, and its identification as 2′-*O*-methyladenylyl-2′-*O*-methyluridylylguanylate (Trim, A. R. & Parker, J. E.) 257–259
Ribosomal ribonucleic acid, *see* Ribonucleic acid, ribosomal
Ribosomal subunits, 60S, binding of, to smooth and rough rat liver endoplasmic-reticulum membranes (Rolleston, F. S. & Mak, D.) 851–853
Ribosomes, loss of protein-synthesizing activity of, *in vitro* associated with loss of viability of rye grains (Roberts, B. E., Payne, P. I. & Osborne, D. J.) 275–286
Root nodules, soya-bean, kinetic properties of the nitrogenous complex from bacteroids of (Bergersen, F. J. & Turner, G. L.) 61–75
Rye (*Secale cereale*) grains, loss of activity of protein-synthesizing systems *in vitro* associated with loss of viability of (Roberts, B. E., Payne, P. I. & Osborne, D. J.) 275–286

Sanfilippo's syndrome, quantitative determination with Alcian Blue 8GX glycosaminoglycans in the urine of human patients with (Whiteman, P.) 351–357
Sclerotinia fructigena, partial purification and properties of β-*N*-acetylglucosaminidase from (Reyes, F. & Byrde, R. J. W.) 381–388
Sea-cucumber (*Holothuria forskali*), isolation of a new neuraminic acid derivative and three types of glycopeptides from the Cuvierian tubules of (Isemura, M., Zahn, R. K. & Schmid, K.) 509–521
Secale cereale, *see* Rye
Serum, human, inhibition by α_2-macroglobulin and other proteins from, of the activity of human liver cathepsin B1 (Starkey, P. M. & Barrett, A. J.) 823–831
Skeletal muscle, *see* Muscle, skeletal
Skin, calf, effect of alcohols and neutral salt on the thermal stability of soluble and precipitated acid-soluble collagen from (Russeli, A. E.) 335–342
Skin, ox, age-related changes in the reducible cross-links of intact collagen fibres from (Robins, S. P., Shimokomaki, M. & Bailey, A. J.) 771–780
Small intestine, *see* Intestine, small
Sodium sulphite, dissociation of human immunoglobulin M by treatment with (Ricardo, M. J., Jr. & Inman, F. P.) 677–682
L-Sorbose, inhibition by D-glucose and its analogues of the penetration of, into human erythrocytes (Barnett, J. E. G., Holman, G. D. & Munday, K. A.) 211–221
Soya bean (*Glycine max*), kinetic properties of the nitrogenase complex from root-nodule bacteroids of (Bergersen, F. J. & Turner, G. L.) 61–75
Spinacea oleracea, *see* Spinach
Spinach (*Spinacea oleracea*), amino acid sequence of cytochrome *c* from (Brown, R. H., Richardson, M., Scogin, R. & Boulter, D.) 253–256
Spleen, horse, molecular weight of the oligomer of apoferritin from, and its stability to dissociation by dilution (Crichton, R. R., Eason, R., Barclay, A. & Bryce, C. F. A.) 855–857
Spleen, human, activity of cathepsin A in (Bowen, D. M. & Davison, A. N.) 417–419
Spleen, human and horse, organ-specificity of ferritin in, and liver (Crichton, R. R., Millar, J. A., Cumming, R. L. C. & Bryce, C. F. A.) 51–59
Spleen, human, reconstitution of glucocerebrosidase activity from two macromolecular components from (Ho, M. W., O'Brien, J. S., Radin, N. S. & Erickson, J. S.) 173–176
Stomach, mouse, properties of the uridine diphosphate glucuronyltransferase activity synthesizing thio-β-D-glucuronides in preparations of (Illing, H. P. A. & Dutton, G. J.) 139–147
Streptococcus faecium A.C.T.C. 9790, selective extraction of teichuronic acids from the cell walls of, and other Gram-positive bacteria (Pavlik, J. G. & Rogers, H. J.) 619–621
Streptomyces strains, peptide inhibitors of DD-carboxypeptidases from (Nieto, M., Perkins, H. R., Leyh-Bouille, M., Frère, J.-M. & Ghuysen, J.-M.) 163–171
Streptomyces strains, specificity of DD-carboxypeptidases from, acting as peptidyltransferases for amino compounds as carboxyl acceptors (Perkins, H. R., Nieto, M., Frère, J.-M., Leyh-Bouille, M. & Ghuysen, J.-M.) 707–718
Succinate, inhibition by, of maize-seedling phosphoenolpyruvate carboxylase (Wong, K. F. & Davies, D. D.) 451–458
Sugars, structural requirements for the binding of, to the sugar-transport system of human erythrocytes (Barnett, J. E. G., Holman, G. D. & Munday, K. A.) 211–221

Sulphite, sodium, dissociation of human immunoglobulin M by treatment with (Ricardo, M. J., Jr. & Inman, F. P.) 677–682
Sulphydryl, *see* Thiol
Supernatant fraction, *see* Cytosol

L-Tartrate, inhibition by, of maize-seedling phosphoenolpyruvate carboxylase (Wong, K. F. & Davies, D. D.) 451–458
Tay–Sachs disease, immunological properties of liver *N*-acetylglucosaminidase from normal humans and from patients with (Carroll, M. & Robinson, D.) 91–96
Teeth, ox, developing, purification and partial characterization of amelogenins from the dental enamel of (Eggert, F. M., Allen, G. A. & Burgess, R. C.) 471–484
Teichuronic acids, selective extraction of, from the cell walls of Gram-positive bacteria (Pavlik, J. G. & Rogers, H. J.) 619–621
Tendon, ox, age-related changes in the reducible cross-links of intact collagen fibres from (Robins, S. P., Shimokomaki, M. & Bailey, A. J.) 771–780
Testis, rat, partial reactions of inositol cyclase from (Barnett, J. E. G., Rasheed, A. & Corina, D. L.) 21–30
Thio-β-D-glucuronides, properties of the uridine diphosphate glucuronyltransferase activity synthesizing, in preparations from various animal sources (Illing, H. P. A. & Dutton, G. J.) 139–147
Thiol groups, protein, use of 3-bromopropionic acid for the determination of (Bradbury, A. F. & Smyth, D. G.) 637–642
Thiols, conversion of the type O (oxidase) forms of cow's milk xanthine oxidase into the type D (dehydrogenase) form in the presence of (Battelli, M. G., Lorenzoni, E. & Stirpe, F.) 191–198
Thrombin, conversion of ox plasma prothrombin into, catalysed by Factor X*a* (Jesty, J. & Esnouf, M. P.) 791–797
Thrombin, ox, determination of the operational molarity of solutions of, by spectrofluorimetric titration (Jameson, G. W., Roberts, D. V., Adams, R. W., Kyle, W. S. A. & Elmore, D. T.) 107–117
Thrombin, release by, of phospholipase A_1 by human blood platelets (Smith, J. B., Silver, M. J. & Webster, G. R.) 615–618
Thymus gland, calf and rabbit, properties of polynucleotides associated with preparations of histones from, and various other sources (Greenaway, P. J. & Murray, K.) 585–591
Thymus gland, calf, interaction of quinacrine mustard with histones from (Selander, R.-K. & de la Chapelle, A.) 757–764
Thymus gland, rat, characterization of a non-histone preparation isolated from histone F1 preparations from, and other sources (Smith, J. A. & Stocken, L. A.) 859–861
Tobacco-mosaic virus, reactivity towards methyl picolinimidate of additional lysine residues in the coat protein of mutants of, and the reagent's usefulness as a probe for investigating the topography of the virus (Perham, R. N.) 119–126
Transpeptidation, specificity of *Streptomyces* DD-carboxypeptidases for amino compounds as carboxyl acceptors for (Perkins, H. R., Nieto, M., Frère, J.-M., Leyh-Bouille, M. & Ghuysen, J.-M.) 707–718
p-(*NNN*-Trimethylammonium)cinnamic acid, determination of the operational molarity of solutions of serine proteinases by spectrofluorimetric titration with the 4-methylumbelliferyl ester of (Jameson, G. W., Roberts, D. V., Adams, R. W., Kyle, W. S. A. & Elmore, D. T.) 107–117
Trinucleotide, alkali-stable, preparation of, from yeast ribonucleic acid and its identification as 2′-*O*-methyladenylyl-2′-*O*-methyluridylylguanylate (Trim, A. R. & Parker, J. E.) 257–259
Triticum vulgare, *see* Wheat
Troponin complex, skeletal-muscle, rabbit, phosphorylation of the '37000 component' (troponin-T) from (Perry, S. V. & Cole, H. A.) 425–428
Troponin-T ('37000 component') phosphorylation of, from rabbit skeletal-muscle troponin complex (Perry, S. V. & Cole, H. A.) 425–428
Trypsin, effects of ligands on the inactivation of wheat-germ aspartate carbamoyltransferase by (Yon, R. J.) 699–706
Trypsin, ox, determination of the operational molarity of solutions of, by spectrofluorimetric titration (Jameson, G. W., Roberts, D. V., Adams, R. W., Kyle, W. S. A. & Elmore, D. T.) 107–117
Tryptophan 5-hydroxylase, partial purification and properties of, from rat small intestine (Noguchi, T., Nishino, M. & Kido, R.) 375–380

Uridine diphosphate 6-deoxygalactose, competitive inhibition and substrate activity of, for uridine diphosphate galactose 4-epimerase from *Escherichia coli* K12 (Spencer, M., Blackburn, P., Ferdinand, W. & Blackburn, G. M.) 421–423
Uridine diphosphate fucose, competitive inhibition and substrate activity of, for uridine diphosphate galactose 4-epimerase from *Escherichia coli* K12 (Spencer, M., Blackburn, P., Ferdinand, W. & Blackburn, G. M.) 421–423
Uridine diphosphate galactose 4-epimerase, competitive inhibition and substrate activity of uridine diphosphate 6-deoxygalactose for, from *Escherichia coli* K12 (Spencer, M., Blackburn, P., Ferdinand, W. & Blackburn, G. M.) 421–423
Uridine diphosphate glucuronyltransferase, properties of the activity of, synthesizing thio-β-D-glucuronides in preparations from various animal sources (Illing, H. P. A. & Dutton, G. J.) 139–147
Uridine monophosphate, effect of, on the inactivation of wheat-germ aspartate carbamoyltransferase by trypsin and denaturing agents (Yon, R. J.) 699–706
Urine, quantitative determination of glycosaminoglycans in, with Alcian Blue 8GX (Whiteman, P.) 351–357
Uterus, rat, immature, oestrogen-stimulated biosynthesis of heterogeneous nuclear ribonucleic acid in (Knowler, J. T. & Smellie, R. M. S.) 689–697

Venom, Malayan-pit-viper, isolation and properties of the coagulant enzyme Arvin from (Hatton, M. W. C.) 799–807
Virus, tobacco-mosaic, *see* Tobacco-mosaic virus
Vitamin C, *see* Ascorbate

Wall, cell, *see* Cell wall
Wheat (*Triticum vulgare*) germ, effects of ligands on the inactivation by trypsin and denaturing agents of aspartate carbamoyltransferase from (Yon, R. J.) 699–706
Wheat (*Triticum vulgare*) germ, purification, composition and specificity of agglutinin from (Allen, A. K., Neuberger, A. & Sharon, N.) 155–162

Xanthine oxidase, characterization of alternative molecular forms of, from mouse liver and duodenum (Duke, E. J., Joyce, P. & Ryan, J. P.) 187–190
Xanthine oxidase, milk, cow, purification, interconversion and properties of the type D (dehydrogenase) and type O (oxidase) forms of (Battelli, M. G., Lorenzoni, E. & Stirpe, F.) 191–198

Yeast, activity of glucose 6-phosphate dehydrogenase from, towards C-2-substituted derivatives of D-glucose 6-phosphate (Bessell, E. M. & Thomas, P.) 83–89
Yeast, activity of glucose phosphate isomerase from, towards deoxyfluoro-D-glucose 6-phosphates (Bessell, E. M. & Thomas, P.) 77–82
Yeast, baker's, kinetics and mechanism of the reaction catalysed by alcohol dehydrogenase from, with a variety of substrates (Dickinson, F. M. & Monger, G. P.) 261–270
Yeast, preparation of an alkali-stable trinucleotide from ribonucleic acid from, and its identification as 2′-*O*-methyladenylyl-2′-*O*-methyluridylylguanylate (Trim, A. R. & Parker, J. E.) 257–259
Yeast, properties of polynucleotides associated with preparations of histones from, and various other sources (Greenaway, P. J. & Murray, K.) 585–591
Yeast, protein kinase activity in commercially available crystalline preparations of alcohol dehydrogenase from (Kemp, B. E., Froscio, M. & Murray, A. W.) 271–274

Zea mays, *see* Maize

rogenase
s of D-
omas, P.)

se from,
(Bessell,

reaction
, with a
Monger,

tide from
ation as
guanylate

ted with
us other
585–591
lly avail-
dehydro-
Murray,